



Chronic Myeloid Leukaemia (CML) Clinical Trials Portfolio





Recruiting CML Trials	
OPTIC	A Randomized, Open-label, Phase 2 Trial of Ponatinib in Patients With Resistant Chronic Phase (CP) CML to Characterize the Efficacy and Safety of a Range of Doses
(Interventional)	Eligibility Criteria:
PI: Prof Adam Mead	Age 18 or over, CML-CP and have received at least two prior tyrosine-kinase inhibitor (TKI) therapies and have demonstrated resistance to treatment OR have documented history of presence of T3151 mutation after receiving any number of prior TKI, ECOG performance status ≤ 2 with adequate end organ function
	A Phase 3, Multi-center, Open-label, Randomized Study of Oral ABL001 Versus Bosutinib in Patients With CML-CP, Previously Treated With 2 or More Tyrosine Kinase Inhibitors
CABL001 A2301	Eligibility Criteria:
(Interventional) PI: Prof Adam Mead	Age 18 or over with a diagnosis of CML-CP, < 15% blasts in peripheral blood and bone marrow, < 30% blasts plus promyelocytes in peripheral blood and bone marrow, < 20% basophils in the peripheral blood, ≥ 50 x 109/L platelets, No evidence of extramedullary leukemic involvement with the exception of hepatosplenomegaly, BCR-ABL ratio ≥ 1% IS, Prior treatment with a minimum of 2 prior ATP-binding site TKIs, Failure or intolerance to the last previous TKI therapy
	A phase 2, multi-center, open-label, randomized study of Oral asciminib added to Imatinib versus continued imatinib versus switch to nilotinib in patients with CML-CP who have been previously treated with imatinib and have not achieved deep molecular response
CABL001 E2201	Eligibility Criteria:
(Interventional) PI: Prof Adam Mead	Age 18 or over with a confirmed diagnosis of CML-CP (< 15% blasts in peripheral blood and bone marrow, < 30% blasts plus promyelocytes in peripheral blood and bone marrow, < 20% basophils in the peripheral blood, $\geq 100 \times 10^9 / L$ platelets), $\geq 1.5 \times 10^9 / L$ neutrophils, $\geq 75 \times 10^9 / L$ platelets, ≥ 9 g/dl haemoglobin, with an adequate end organ function, No evidence of extramedullary leukemic involvement with the exception of hepatosplenomegaly, ≥ 24 months of Imatinib treatment, BCR-ABL ratio > 0.01% IS and $\leq 1\%$ IS
INForMeD	An observational and biological research study to investigate the genetic and cellular basis of sporadic and familial myeloid disorders
(Non-interventional)	Eligibility Criteria:
PI: Prof Adam Mead	Age 2 or over, Patients under investigation for or diagnosed with a myeloid or related disorder, Patient willing to give consent to the study

Haematology Research contact details	
Prof Adam Mead	Email: adam.mead@imm.ox.ac.uk Tel: 01865 222324
Research Nurse Team	Email: LP.haematology.RN@oxnet.nhs.uk

Last updated: March 2019