

Department of Clinical Haematology

Prevention and Management of Tumour Lysis Syndrome (TLS) in Adults

Background

Tumour Lysis Syndrome (TLS) is a metabolic complication that can occur during systemic anti-cancer treatment (SACT) or radiotherapy in certain haematological malignancies and very occasionally solid tumours (such as germ cell tumours or bulky sarcomas).

TLS is characterised by the rapid development of two or more of the following:

- **Hyperuricaemia**
- **Hyperkalaemia**
- **Hyperphosphataemia**
- **Hypocalcaemia**
- **Clinical TLS** (biochemical signs alongside the following clinical manifestations):
 - Acute kidney injury (rising creatinine)
 - Cardiac arrhythmia,
 - Seizure or sudden death.

Acute release of intracellular products (e.g., urate, phosphate, potassium) into the circulation is a consequence of rapid cell lysis from radiosensitive or SACT sensitive rapidly proliferating cells.

Hypocalcaemia occurs as result of precipitation of calcium phosphate in soft tissues due to the acute development of hyperphosphataemia.

Acute renal insufficiency results from the precipitation of uric acid (common pre-SACT) and/or calcium phosphate crystals in renal tubules (common following SACT).

Risk Factors

There are several well recognised risk factors for the development of laboratory and clinical TLS, including:

- High tumour burden
- High grade tumours with rapid cell turnover
- Pre-existing renal impairment or renal involvement by tumour
- Increased age
- Treatment with highly active, cell-cycle specific agents
- Concomitant use of drugs that increase uric acid levels, for example: alcohol, ascorbic acid, aspirin, caffeine, cisplatin, diazoxide, thiazide diuretics, adrenaline (epinephrine), ethambutol, levodopa, methyldopa, nicotinic acid, pyrazinamide, phenothiazines and theophylline.

Department of Clinical Haematology

Definition

Evidence of **laboratory tumour lysis** syndrome is defined as two or more of the following (Cairo & Bishop, 2004):

uric acid	≥ 476 micromol/L or 25% increase from baseline
potassium	≥ 6mmol/L or 25% increase from baseline
calcium	≤ 1.75mmol/L or 25% decrease from baseline
phosphate	≥ 1.45 mmol/L or 25% increase from baseline

Additional investigations:

- LDH (elevated)
- Serum creatinine >1.5 times the upper limit of normal (or patient baseline)
- Serum urea (elevated)
- Urine pH <5
- ECG - arrhythmia

Risk Stratification for TLS

Risk stratification enables the identification of patients that should be offered active monitoring, hydration and rasburicase prophylaxis, as opposed to the low and intermediate risk groups, in which hydration +/- allopurinol prophylaxis should suffice (consider inpatient monitoring in borderline situations).

High-risk patients can be identified by the following criteria:

- Burkitt's or Burkitt's-like Lymphoma
- Lymphoblastic Lymphoma
- ALL WCC > 100 x 10⁹/l
- AML WCC > 100 x 10⁹/l
- CML in blast crisis WCC > 100 x 10⁹/l
- Or WCC < 100 and LDH > 2 times upper limit of normal
- High-grade Lymphoma with bulky disease – defined by LDH more than twice the upper limit of normal or tumour bulk >10 cm in diameter.

Note:

Diseases which are traditionally low risk (e.g., CLL) may become very high risk with the use of novel therapeutics. Special care should be taken in these therapy-specific circumstances, irrespective of pathology.

Increasing age, pre-existing renal impairment and renal involvement by tumor are also factors which increase risk and may push the treating clinician to consider specific patients as high-risk.

Department of Clinical Haematology

Prevention of Tumour Lysis Syndrome

1. Low/intermediate risk patients (prophylaxis)

Low/intermediate risk patients can be managed with a combination of hydration and allopurinol.

Hydration	<ul style="list-style-type: none"> Start hydration before anti-cancer treatment. Aim for total hydration fluid of 2-3L in 24 hours. AVOID additional potassium in hydration fluids
Medication	<ul style="list-style-type: none"> Allopurinol 300mg ONCE daily for 7 days*, start 24 - 48 hours prior to SACT where possible. Reduce allopurinol to 100 - 200mg ONCE daily where creatinine clearance (CrCl) is <20ml/min.

* Allopurinol dosing follows standard and established practice within haematology; however, the adult dosing schedule allows 200 – 400 mg/m²/day in 1 or 3 divided doses, up to a maximum of 800 mg daily.

Observations:

- Daily weight
- Vital signs
- Fluid balance
- Blood sampling (creatinine, urea & electrolytes; including calcium, phosphate, potassium and magnesium, uric acid, LDH) in addition to usual SACT monitoring.

Additional considerations:

- Minimise use of concurrent nephrotoxic drugs, avoid thiazide or potassium-sparing diuretics.
- A select group of intermediate-risk group patients may warrant inpatient monitoring with enhanced hydration and electrolyte monitoring.
- Patients allergic to allopurinol should be considered for rasburicase.

Deterioration during prophylactic treatment

In the case of deteriorating biochemical, or clinical markers, there should be prompt initiation of rasburicase.

In individuals contra-indicated for rasburicase, allopurinol doses can be increased up to a maximum of 800mg/day (in 2-3 divided doses). Early liaison with ITU/renal dialysis is essential as allopurinol will not prevent further renal decline.

2. High-risk patients (prophylaxis)

High risk patients should be managed with a combination of IV hydration and rasburicase.

Hydration	<p>IV hydration should start 2 days before anti-cancer treatment. Administer IV fluids to maintain urine output of 100ml/m²/hour Consider loop diuretic to force diuresis if not adequate.</p> <p>AVOID additional potassium in hydration fluids</p>
------------------	---

Department of Clinical Haematology

Medication	<p>Rasburicase 3mg IV stat 30-60mins prior to anti-cancer treatment. A second dose can be considered within individual cases.</p> <p>For G6PD deficient individuals: Allopurinol 200 – 400mg/m²/day in 1-3 divided doses, up to a maximum of 800mg daily.</p>
-------------------	--

Rasburicase prophylaxis dose

The licensed dose is 0.2mg/kg/day IV for up to 7 days (Sanofi, 2021), however a meta-analysis reviewing the effectiveness of a single lower fixed dose of rasburicase in high-risk patients concluded non-inferior clinical benefit to the licensed dose for high-risk prophylaxis (Feng et al, 2013).

British Society of Haematology (BSH) recommends a fixed dosage of 3mg in the prophylactic setting (Jones et al, 2015) which has since been adopted by several Trusts across the UK.

We endorse the recommendation for a **single, fixed dose of 3mg rasburicase** for high-risk patients prior to SACT, with regular monitoring and **application of a second dose if required**.

Observations:

- Daily weight
- Vital signs
- Fluid balance
- Blood sampling (creatinine, urea & electrolytes; including calcium, phosphate, potassium and magnesium, uric acid, LDH) in addition to usual anti-cancer treatment monitoring.
 - OUH out of hours: A venous blood gas must be taken and tested in parallel to standard laboratory bloods as there is no on site out of hours biochemistry lab at the Churchill. The standard bloods should be couriered the John Radcliffe Hospital urgently.

Additional considerations:

- Minimise the use of concurrent nephrotoxic drugs, avoid thiazide or potassium-sparing diuretics.
- Several contra-indications to rasburicase treatment exist, please refer to Appendix 1.

Deterioration during prophylactic treatment

In the case of deteriorating biochemical, or clinical markers, there should be prompt escalation to the full protocol for management of established TLS.

Department of Clinical Haematology

Management of Established Tumour Lysis Syndrome

1. Liaise with ITU/ renal unit as early as possible.

2. Hydration

- Ensure IV fluids are running at greater than 3 litres/m²/day (i.e., twice usual maintenance)
- It is critical that potassium is **NOT** added to the hydration fluid.
- Aim for urine output greater than 100ml/ m²/ hr. Monitor urine output hourly.
- Consider loop diuretic to force diuresis if not adequate.
- Biochemistry every 6 hours or the first 24 hours after diagnosis
- Maintain strict fluid balance chart and place urinary catheter.
- Assess fluid balance formally every 4 hours.
- Weigh twice daily.
- Fluid retention may be treated with IV furosemide (0.5- 1mg/kg) or mannitol (0.5mg/kg) if weight gain is > 3 kg. In the event of severe oliguria or anuria, a single dose of furosemide (2-4mg/kg) may be considered to improve or initiate urinary output.

3. Rasburicase

- **Initiate rasburicase at 0.2 mg/kg IV for 3 - 7 days** depending on clinical and biochemical parameter. No dose adjustment is necessary in renal or hepatic impairment.
- **Where rasburicase is contra-indicated** - ensure maximum dose of allopurinol is administered (up to max 800mg daily in 2 - 3 divided doses). Early discussions with ITU/renal dialysis are essential.
- Refer to Appendix 1 for rasburicase contra-indications, side-effects, reconstitution, and administration details.

4. Patient Monitoring

- A patient with established TLS can deteriorate rapidly, and close monitoring is essential to anticipate escalations in care.
- Clinical monitoring:
 - Vital sign monitoring (SaO₂, RR, HR, BP and temperature) at least 4 to 6-hourly.
 - Urine output monitoring on an hourly basis
 - Weight, at least daily
 - Clinical examination of fluid status at least 6-hrly
 - ECG monitoring at baseline, and as indicated
- Blood monitoring:
 - Check potassium, phosphate, calcium, magnesium, urea, creatinine and urate at least 6-hrly initially, using blood gas samples in parallel to acquire rapid potassium values.
 - Blood samples and rasburicase: Following administration of rasburicase, uric acid levels continue to drop in vitro, and so urate results will be spuriously low making interpretation difficult. This effect can be mitigated, though not completely avoided, by immediately placing the sample on ice and rushing it to the biochemistry laboratory for rapid assay. This is not always feasible in day-to-day practice; in which case the biochemical and clinical picture should be assessed in its entirety to guide

Department of Clinical Haematology

management.

- If there is no evidence of TLS reduce frequency of sampling.

5. Treatment of hyperkalaemia

- Clinical manifestations include nausea, anorexia, vomiting, diarrhoea, neuromuscular and cardiac abnormalities.
- Treat according to local guideline for management of hyperkalemia
- Information re ECG monitoring etc. is contained within the OUH link immediately above, print out the document for pathway management.

6. Treatment of hyperphosphataemia

- Clinical manifestations include nausea, vomiting, diarrhoea, lethargy and seizures.
- If hydration and timely administration of rasburicase do not prevent significant hyperphosphataemia, it can be hard to control phosphate levels other than by dialysis.
- The temporary use of aluminum hydroxide 50–150 mg/kg/day has been described but is slow to act and poorly tolerated, thus is not routinely recommended in this setting.
- Avoid calcium supplements except in neuromuscular irritability.

7. Treatment of hypocalcaemia

- Clinical manifestations include muscular (cramps and spasms, paraesthesias, tetany), cardiovascular (ventricular arrhythmias, heart block, hypotension, prolonged QT interval) and neurological complications (confusion, delirium, hallucinations, and seizures)
- **Treatment of asymptomatic hypocalcaemia is not recommended** as the risk of precipitating metastatic calcification is high, especially in the setting of hyperphosphataemia.
- Symptomatic hypocalcaemia should be treated with IV calcium gluconate as per local guideline. The aim of treatment is to treat the symptoms but not to normalize the biochemical parameters.

Sodium bicarbonate: Alkalinisation of urine using sodium bicarbonate to raise the urine pH is no longer recommended.

8. Indications for haemodialysis/intensive care

- Potassium ≥ 6.5 mmol/L at 4 hours despite interventions.
- Rising urea, creatinine, or phosphate despite interventions above.
- Central nervous system toxicity secondary to uraemia.
- Metabolic acidosis
- Fluid overload unresponsive to diuretics

Department of Clinical Haematology

References

1. Cairo M.S. & Bishop M. (2004) Tumour lysis syndrome: new therapeutic strategies and classification. British Journal Haematology, 127, 3–11.
2. Cairo M.S., Coiffier, B., Reiter, A. & Younes, A. (2010) Recommendations for the evaluation of risk and prophylaxis of tumour lysis syndrome (TLS) in adults and children with malignant diseases: an expert TLS panel consensus. British Journal of Haematology, 149, 578–586
3. Feng X., Dong, K., Pham, D., Pence, S., Inciardi, J. & Bhutada, N.S. (2013) Efficacy and cost of single dose rasburicase in prevention and treatment of adult tumour lysis syndrome: a meta-analysis. Journal of Clinical Pharmacy and Therapeutics, 38, 301–308.
4. Jones GL, Will A, Jackson GH, Webb NJ, Rule S; British Committee for Standards in Haematology. Guidelines for the management of tumour lysis syndrome in adults and children with haematological malignancies on behalf of the British Committee for Standards in Haematology. Br J Haematol. 2015 Jun;169(5):661-71
5. Oxford University Hospitals NHS Trust. Guidelines for the management of hyperkalemia in adults, Sep 2015.
6. Oxford University Hospitals NHS Trust. Guidelines for the management of acute hypocalcaemia in adults, Oct 2014.
7. Sanofi®. Fasturtec 1.5mg/ml power and solvent for concentrate for solution for infusion. [Online] eMC UK Summary of Product Characteristics. 2021. Available from: <https://www.medicines.org.uk/emc/product/1316> (accessed 15/12/2021)

Review

Author: (V 1.0): Claire Humphries, Specialist Pharmacist

Audit: These processes are subject to the OxBMT/IEC audit programme

Circulation: NSSG Haematology Website

Name	Revision	Date	Version	Review date
Dr Graham Collins, Haematologist	Review	Jul 11	2.1	Jul 13
Dr Jaimal Kothari, Haematologist Cheuk-Kie Cheung, Nadjoua Maouche, Specialist Pharmacists. Rachel Miller, Ward Sister. Sandy Hayes, Quality manager	Full review: change in Rasburicase management and sampling audit.	Mar 16	3.0	Dec 16
Faouzi Djebbari, Advanced Haematology Pharmacist	Change in timing of rasburicase administration	Feb 20	3.1	Jun 20
Dr Yisu Gu and Dr Edmund Watson, Haematology SpR. Donna Constantine, Advanced Haematology Pharmacist Denise Wareham, BMT Senior Specialist Nurse	BMJ Best Practice updates; Rasburicase patient monitoring ; out of hours access to rasburicase; G6PD high risk populations; Additional minor amendments	Dec 21	3.2	Dec 23
Donna Constantine, Advanced Cancer Pharmacist Lymphoma NSSG Review Meeting	Rasburicase prophylactic dosing recommended as 3mg fixed dose as per BSH guidance.	Jul 2023	4.0	Dec 2024
Unknown author Donna Constantine, Advanced Cancer Pharmacist	Minor clarification re: out of hours blood sampling specific to OUH. Formatting.	May 2024	4.1	Jul 2026

Department of Clinical Haematology

Appendix 1- Rasburicase administration, contra-indications, and side-effects

Presentation

Rasburicase 1.5mg in 1ml vial

Rasburicase 7.5mg in 5ml vial

Dosing

Prophylaxis of TLS (Off-label use):

3mg IV as a single dose 30 – 60mins prior to anti-cancer treatment or high-dose steroids. A 3mg dose can be repeated once more in individual circumstances.

Treatment of TLS:

0.2mg/kg IV ONCE daily for 3 – 7 days as necessary

Round all doses to the nearest 1.5mg vial.

Side effects

Very common ($\geq 1/10$): Headache, diarrhoea, vomiting, nausea, fever

Common ($\geq 1/100$ - $< 1/10$): Allergic reaction (rash, urticaria)

Uncommon ($\geq 1/1,000$ to $< 1/100$): Haemolysis, haemolytic anaemia, hypotension, bronchospasm, methaemoglobinaemia, severe hypersensitivity

Rare: Anaphylaxis and involuntary muscle contraction has been reported.

Contra-indications

Rasburicase treatment is contraindicated in:

- Allergy to rasburicase, or excipients
- Glucose 6-phosphate dehydrogenase (G6PD) deficiency
- History of haemolytic anaemia, or presence of other cellular disorders known to cause haemolytic anaemia.
- Methaemoglobinaemia

G6PD deficiency is common in individuals originating from Africa, Asia, the Mediterranean region, and the Middle East; it can also occur less frequently in all other individuals. G6PD deficiency is more common in males than it is in females. A G6PD test is recommended before commencing treatment in individuals with strong heritage from or originating from these areas. G6PD test results may not be immediately available upon request and so should be requested pro-actively during patient diagnostic work-up.

Reconstitution and administration

- Medication is stored within the fridge.
- Reconstitute each vial of rasburicase required with the vial of solvent provided. Mix by swirling gently. Do not shake.
- Dilute in 50ml sodium chloride 0.9% and infuse over 30 minutes.
- The diluted drug can be given peripherally.

Supply out of hours (OUH Specific)

Department of Clinical Haematology

- Rasburicase vials are stored within the fridge of the Churchill Hospital emergency drug cupboard (EDC), and within the JR EDC fridge for out of hours use.
- Between 6pm – 9pm the CH EDC access key can be obtained from UGI ward. This key will require signing out and you will have to present a Trust ID badge. Between 9pm – 8am contact the Night Nurse Practitioner via Switchboard for access to the EDC.
- Access at the JR Hospital differs and can be accessed via security after pharmacy closes, normally between the hours of 12am – 8am, when the resident is not on site.
- If the EDC stock is missing, please contact the on-call pharmacist via Switchboard.