

Guidelines for Supportive Care in Myeloma

British Committee for Standards in Haematology in conjunction with the UK Myeloma Forum (UKMF)

Address for correspondence:

BCSH Secretary
British Society for Haematology
100 White Lion Street
London
N1 9PF
e-mail bcsh@b-s-h.org.uk

Writing group:

JA Snowden¹, S Ahmedzai², J Ashcroft³, S D'Sa⁴, T Littlewood⁵, E Low⁶, H Lucraft⁷, R Maclean¹, S Feyler⁸, G Pratt⁹, J Bird¹⁰

Disclaimer:

While the advice and information in these guidelines is believed to be true and accurate at the time of going to press, neither the authors, the British Society for Haematology nor the publishers accept any legal responsibility for the content of these guidelines.

Date for initial guideline review: January 2012

1Department of Haematology, Sheffield Teaching Hospitals NHS Trust
2Department of Palliative Medicine, Sheffield Teaching Hospitals NHS Trust
3 Department of Haematology, Leeds Teaching Hospitals NHS Trust
4Department of Haematology, University College, London
5 Department of Haematology, John Radcliffe Hospital, Oxford
6 Myeloma UK, Edinburgh
7 Department Clinical Oncology, Freeman Hospital, Newcastle
8 Department of Haematology, Calderdale and Huddersfield NHS Trust
9 Department of Haematology, Heartlands Hospital, Birmingham
10 Avon Haematology Unit, Bristol Haematology and Oncology Centre, Bristol

CONTENTS

1. Introduction: definitions and methodology
2. Anaemia
3. Haemostasis and thrombosis issues
 - 3.1 *Bleeding risks*
 - 3.2 *Thrombotic risks*
4. Infection
5. Pain management
 - 5.1 *Prevalence and impact of pain in myeloma*
 - 5.2 *Assessment of pain*
 - 5.3 *Interventions*
 - 5.4 *Collaboration with specialist services*
6. Peripheral neuropathy
 - 6.1 *Scope*
 - 6.2 *Causes*
 - 6.3 *Assessment and investigation of peripheral neuropathy*
 - 6.4 *Management of peripheral neuropathy*
7. Control of other symptoms
 - 7.1 *Nausea and vomiting*
 - 7.2 *Loss of appetite (anorexia)*
 - 7.3 *Constipation and diarrhoea*
 - 7.4 *Mucositis*
 - 7.5 *Sedation*
 - 7.6 *Fatigue*
8. Bisphosphonate induced osteonecrosis of the jaw
9. Complementary therapies
 - 9.1 *Definition*
 - 9.2 *The patient's preference*
 - 9.3 *Potential benefits of complementary therapy*
 - 9.4 *Commonly used complementary therapies*

10. End of life care

10.1 Definitions

10.2 Recognition of end of life

10.3 Continuation of anti-cancer and other active therapies

10.4 Last days of life

11. Conclusions – the ‘holistic needs assessment’ and clinical research parameters

12. Declaration of interest

13. Levels of evidence and grades of recommendation

1. Introduction: definitions and methodology

In the vast majority of patients, myeloma is an incurable disease. In recent years, an evolving array of effective treatments have increased the ability to achieve disease control repeatedly to the extent that myeloma now has the potential to be a chronic disease in many patients. The improvement in survival is reflected by the UK MRC myeloma trials and other epidemiological data, and, for example, with modern management younger patients responding to intensive treatment can now expect to survive for a median of approximately 7 years from diagnosis (Morgan *et al*, 2009) or potentially longer (Brenner *et al*, 2008; Brenner *et al*, 2009). However, the improved survival of patients with myeloma has resulted in an increasing symptom burden due not only to the disease itself, but to also the cumulative effects of treatments. The major challenge of modern myeloma management is therefore matching the progress made in prolonging survival through disease control whilst optimising global quality of life with effective supportive care measures from initial diagnosis to end of life care.

The aim of these guidelines is to summarise a national consensus of the haematological community and colleagues involved in the supportive care of patients with myeloma. For the purpose of these guidelines the definition adopted in the UKNHS NICE Guidance (2004b) has been used i.e. “care that helps the patient and their family to cope with cancer and treatment of it – from pre-diagnosis and treatment, to cure, continuing illness or death and into bereavement. It helps the patient to maximise the benefits of treatment and to live as well as possible with the effects of the disease. It is given equal priority alongside diagnosis and treatment.” This definition is sufficiently broad to cover not only to cover symptomatic treatment and palliative care but also the wide range of management options considered to be ‘haematological supportive care’, including anti-infectives, transfusion therapy, anticoagulation and growth factors.

Although much of the supportive care can be provided by the haematologists and their teams principally responsible for the care of patients with myeloma, if patients fail to respond or experience intolerable side-effects, advice should be sought from other specialist teams. In some patients, satisfactory symptomatic management is often only achieved through good multidisciplinary collaboration and specialist input from colleagues in Palliative Medicine, Pain Management, Clinical Oncology and Orthopaedics. This is

probably best achieved through the routine MDT meetings (2003b). Management of psychological aspects is also important. Outside of the hospital environment referral can be made to the community palliative care team or local hospice service.

These guidelines should be read in conjunction with the British Committee for Standards in Haematology (BCSH) Guidelines for Diagnosis and Management of Multiple Myeloma 2010 (Bird *et al*, submitted) which they complement. Management of symptoms in patients with myeloma at all stages should follow the principles of evidence-based palliative medicine, where possible. Where appropriate, levels of evidence are provided. As there was a view that many haematologists would welcome specific guidance in certain areas, certain therapeutic regimens have been suggested, most of which would be in the remit of haematological practice. However, in other areas, the guidelines make suggestions for specialist care outside of haematology. However, these guidelines should not be taken as prescriptive as variations exist. The guidance may not be appropriate to all patients and individual patient circumstances or clinician preferences may reasonably favour an alternative approach.

The draft guideline was produced by the writing group consisting of authors, assisted by other members of the broader BCSH Myeloma Guidelines group. Involvement of patient advocacy was achieved through Myeloma UK. The guidelines were subsequently revised by consensus by the UK Myeloma Forum Executive and members of the Haemato-Oncology Task Force of the BCSH. The guidelines were then reviewed by a sounding board of approximately 100 UK haematologists, the BCSH (British Committee for Standards in Haematology) the British Society for Haematology Committee and the comments incorporated where appropriate. The AHCPR system was used to quote levels and grades of evidence where specified which is outlined in section 13 and on the BCSH website (www.bcsguidelines.com) In preparing these guidelines the authors have considered overall cost-effectiveness of recommended interventions as well as clinical efficacy data but formal health economic assessments have not been carried out.

The use of these guidelines to assist management of individual patients should be combined with appropriate professional training. All drug doses should be checked at reference sources, such as the British National Formulary (BNF), Palliative Care Formulary (PCF) or similar. The authors, BCSH or publishers cannot take legal responsibility for individual patient management.

2. Management of anaemia

Anaemia (haemoglobin concentration $<12.0\text{g/dl}$) is common in myeloma. At diagnosis it is present in approximately 75% of patients (Kyle *et al*, 2003). In most patients the anaemia will be normochromic and normocytic and attributed to the myeloma itself and/or the myelosuppressive effect of the chemotherapy. Other causes such as haematinic deficiency or bleeding should be excluded. Fatigue is also reported by many patients and may be caused by both physical and psychological factors related to the disease and its treatment but anaemia has been shown to be an important contributory factor (Cella *et al*, 2004). A European wide survey in patients with myeloma suggested that prevalence of anaemia during chemotherapy is around 85% (Birgegard *et al*, 2006).

Anaemia may be managed by blood transfusion or treatment with erythropoietic stimulating agents (ESA). Blood transfusion may be very helpful in the short-term correction of moderate to severe anaemia in a symptomatic individual. Minimally or asymptomatic individuals with mild or moderate anaemia (due to their disease) may be observed and some will become less anaemic as the myeloma is controlled with chemotherapy. ESA treatment is recommended for anaemic patients with myeloma with associated renal impairment (Locatelli *et al*, 2004). ESA doses of $< 20,000\text{iu/week}$ may be adequate in patients where renal disease is the main cause of the anaemia. In the UK, it may be necessary to refer the patient to a renal physician to access NHS funding for ESAs.

Data from randomised trials, which have included patients with myeloma, suggest that ESA increase the haemoglobin concentration in around two thirds of patients, reduce transfusion need and have a significant positive impact on quality of life (Hedenus *et al*, 2003; Littlewood *et al*, 2001; Osterborg *et al*, 2002).

Recent guidelines from the American Society of Haematology and American Society of Clinical Oncology recommend ESA in symptomatic patients who are receiving concurrent chemotherapy with a Hb concentration which is approaching or fallen below 10g/dl (Rizzo *et al*, 2008). The target haemoglobin concentration should be around 12.0g/dl . European guidelines suggest consideration is given to initiating ESA in symptomatic patients with a

Hb of < 11.0g/dl (Bokemeyer *et al*, 2007). In the UK, The National Institute of Clinical Excellence has not recommended treatment for cancer-treatment related anaemia (except in ovarian cancer).

In patients undergoing high dose therapy, treatment with ESA during the period of cytopenia have no impact on reducing transfusion need. In contrast some studies with small numbers of patients suggest that starting treatment with ESA on day +30 can increase haemoglobin concentration and reduce transfusion need after both autologous and allogeneic transplants (Baron *et al*, 2003; Vanstraelen *et al*, 2005).

There is an increased risk of thrombotic events in patients with cancer who are treated with ESAs (relative risk 1.67). Concerns about an increased mortality in ESA treated patients have also been raised but there are no data suggesting that the outcome of patients with myeloma is worse after ESA treatment. In patients with cancer being treated with chemotherapy there is no difference in survival in randomised trials between those receiving an ESA and those treated by placebo (relative risk 1.04) (Bohlius *et al*, 2009).

Myeloma patients being treated with thalidomide or lenalidomide have an increased risk of thrombotic events. There are conflicting reports about whether this risk is further enhanced by treatment with ESA (Bennett *et al*, 2006; Menon *et al*, 2008).

Recommendations

- **A therapeutic trial of ESA should be considered in a patient with persistent symptomatic anaemia (typically haemoglobin concentration <10.0 g/dl) in whom haematinic deficiency has been excluded (Grade A recommendation; level 1b evidence)**
- **One of Darbepoetin (6.25micrograms/kg every 3 weeks), Epoetin alfa (40,000iu once weekly or Epoetin Beta (30,000iu once weekly) sub-cutaneously can be chosen. Dose doubling after 4 weeks in patients with a haemoglobin increase of <1.0g/dl can be considered (Grade B recommendation; level 2a evidence). ESA treatment should be stopped after 6-8 weeks if there has been no haemoglobin response. The haemoglobin concentration should not rise above 12.0g/dl (Grade C recommendation; level IV evidence).**
- **Patients with anaemia attributed to renal failure should have a trial of treatment with ESA (Grade B recommendation; level IIa evidence)**

- **True or functional iron deficiency occurring during treatment with an ESA should be treated with intravenous iron (NCCN guideline 2010).**

3. Haemostasis and thrombosis issues

3.1 Bleeding risks

Bleeding is rarely associated with myeloma at presentation, more commonly resulting from disease progression, thrombocytopenia (immune mediated or due to marrow infiltration), renal failure, infection and therapy toxicity. Paraproteinaemia in myeloma has, however, been reported to cause bleeding due to acquired von Willebrand's disease (Sampson *et al*, 1983), platelet dysfunction (Di Minno *et al*, 1986), fibrin polymerisation defects (Coleman *et al*, 1972), hyperfibrinolysis (Sane *et al*, 1989) or circulating heparin-like anticoagulants (Llamas *et al*, 2001) and, in AL amyloidosis, factor X deficiency (Mumford *et al*, 2000). Generally, a reduction in plasma paraprotein (by plasmapheresis and/or cytoreductive therapy) will improve haemorrhagic manifestations due to, or exacerbated by, paraproteinaemia. Prothrombin complex concentrates, recombinant factor VIIa and splenectomy have been reported to be successful in the management of some bleeding episodes in patients with factor X deficiency due to AL amyloidosis (Boggio and Green 2001; Thompson *et al*, 2010). There is no consensus on the management of acquired von Willebrand's disease in myeloma. Bleeding episodes have been managed with variable success with desmopressin (transient if any response), intravenous immunoglobulin infusions (usually result in both clinical and laboratory improvement), factor VIII/ von Willebrand's factor concentrates (variable clinical and laboratory responses)(Federici 2006). Management of bleeding in myeloma patients will need to be individualised and no general recommendation can be made.

3.2 Thrombotic risks

Myeloma and other plasma cell disorders have a well-established association with venous thromboembolism (VTE) (Srkalovic *et al*, 2004). A recent retrospective study of US Veterans Affairs hospital records found the incidence of DVT to be 8.7/1000 in patients with myeloma, in comparison to patients with MGUS (3.1/1000) or those without plasma cell disorders (0.9/1000)(Kristinsson *et al*, 2008). Active disease, cancer therapies, infection, previous VTE, immobility and paraplegia are all well-recognised additional risk factors for VTE in hospitalised patients. Thalidomide and lenalidomide have been demonstrated to further increase this risk. The risk of VTE with lenalidomide

alone appears to be lower than with thalidomide. Neither drug used as monotherapy significantly increases risk, but when combined with high-dose steroids or cytotoxic agents the risk of VTE increases significantly (Dimopoulos *et al*, 2007; Richardson *et al*, 2006a; Weber *et al*, 2007). The risk of VTE appears to be higher in patients with newly diagnosed myeloma (NDMM) treated with lenalidomide and dexamethasone (25-75%) (Rajkumar *et al*, 2009; Zonder *et al*, 2006).

3.2.1 Assessment of risk of VTE

As recommended for all patients when admitted to hospital, those with myeloma should be assessed for their risk of developing a VTE, and appropriate chemical thromboprophylaxis (low molecular weight heparin (LMWH) or fondaparinux) given.

All myeloma patients starting thalidomide or lenalidomide should undergo a risk assessment for VTE (see table 1). The reported rates of VTE associated with thalidomide and lenalidomide vary considerably depending on whether the patient is newly diagnosed or relapsed/refractory, and are influenced by the concomitant use of erythropoiesis stimulating agents. This assessment needs to take into consideration patient factors (e.g. previous VTE, obesity, co-morbidities), myeloma related factors (e.g. disease burden, hyperviscosity) and treatment related factors (e.g. concurrent use of high-dose steroids). Patients taking hormone replacement therapy should be encouraged to discontinue it where possible.

Other patients who are *not* receiving thalidomide or lenalidomide may also be at risk of VTE and thromboprophylaxis may be appropriate. This is particularly true of those admitted for management of acute episodes (infection, dehydration, pain management), where their risk of thrombosis may increase dramatically. These patients' need for pharmacological thromboprophylaxis should be considered on a case-by-case basis.

3.2.2 Selection of thromboprophylactic approach

The role of LMWH in preventing VTE in medical and surgical patients is well recognised, although the efficacy of aspirin remains uncertain, and low dose warfarin has generally been shown to be ineffective (Geerts *et al*, 2004). In myeloma, the

evidence base is limited and recommendations combine both broader principles and, where available, specific data.

In myeloma, a number of different thromboprophylactic strategies have been utilised in patients receiving thalidomide or lenalidomide; aspirin, LMWH or warfarin (fixed low dose or adjusted dose to achieve an INR of 2.0-3.0). Both aspirin and LMWH have been reported to reduce the incidence of VTE in patients receiving combination therapies containing thalidomide or lenalidomide. (Zonder *et al*, 2006) found that the incidence of symptomatic VTE with lenalidomide and dexamethasone fell from 75% to 15% with the introduction of aspirin 325mg daily. (Baz *et al*, 2005) also reported a fall in the incidence of symptomatic VTE (from 58% to 18%) with the introduction of aspirin 81mg daily in patients receiving thalidomide with combination chemotherapy. (Palumbo *et al*, 2006) found that 17% of patients receiving MPT developed symptomatic VTE, but with the introduction of enoxaparin thromboprophylaxis, the incidence of symptomatic VTE fell to 3%. (Zangari *et al*, 2004) also demonstrated a fall in the incidence of VTE with the addition of prophylactic LMWH in patients receiving combination chemotherapy in addition to thalidomide. In this study, fixed low dose warfarin did not reduce the risk of symptomatic VTE. Adjusted dose warfarin (INR 2.0-3.0) or therapeutic dose LMWH have also been used in a small study of 26 patients receiving primary induction with thalidomide and dexamethasone with a VTE rate of 8% (Wang *et al*, 2005). There is no evidence of benefit with fixed low-dose warfarin or using target INR less than 2.

As yet, no prospective randomized study comparing thromboprophylaxis strategies in patients receiving thalidomide or lenalidomide-containing treatment regimens has been completed. Despite the lack of clear evidence in this area, the International Myeloma Working Group recently published some recommendations regarding VTE prophylaxis in patients treated with thalidomide or lenalidomide, incorporating both patient-related and treatment-related risk factors for VTE (Palumbo *et al*, 2008c).

Risk factors are categorised into:

- 1) The diagnosis of myeloma itself (as a cancer) and hyperviscosity
- 2) Individual, including previous VTE or inherited thrombophilia, body mass index >30kg/m², central venous catheter in situ, comorbidities (cardiac disease, chronic renal

disease, diabetes, acute infection and immobilisation), surgery, anaesthesia, and trauma, and ESA administration

3) The choice of myeloma therapy, specifically high dose dexamethasone ≥ 480 mg per month, doxorubicin and multi-agent chemotherapy.

Incorporating co-morbidities, the majority of patients undergoing treatment for myeloma with thalidomide or lenalidomide will therefore have ≥ 2 risk factors, including all those receiving high dose dexamethasone or multiagent chemotherapy and, based on this approach, the recommendation is for LMWH (equivalent of enoxaparin 40mg od or dalteparin 5000U od) or adjusted dose warfarin (INR 2.0-3.0). Prophylactic aspirin 81-325mg once daily should only be used in patients with ≤ 1 risk factor. There appears to be no benefit of thromboprophylaxis in the relatively rare situation of patients on thalidomide or lenalidomide monotherapy.

Despite these recommendations, the risks of VTE always need to be individually balanced against the risks of bleeding (e.g. in settings where thrombocytopenia may develop or congenital or acquired haemostatic disorders) and other complications (e.g. osteoporosis). Moreover, safe and optimal administration and monitoring of LMWH and warfarin in a predominantly elderly population may present challenges. Until the results of studies comparing the efficacy of aspirin to LMWH or warfarin in patients receiving such regimens, these recommendations appear sensible (Palumbo *et al*, 2008c). Those on lenalidomide with one additional risk factor may receive aspirin 75mg once daily but there is no evidence that this is protective for a similar group on thalidomide. The latter and those with two additional risk factors require greater protection in the form of prophylactic-dose low molecular weight heparin (LMWH). Those with 3 or more risk factors need full anticoagulation with either treatment-dose LMWH or warfarin with target INR of 2.5.

Thrombocytopenia is a frequent obstacle to safe thromboprophylaxis, related to both marrow infiltration and myeloma treatment. Those starting LMWH should have a baseline platelet count and consideration given to checking further platelet counts every 2 to 4 days for the first 2 weeks of treatment, to screen for heparin-induced thrombocytopenia (HIT) (with a platelet count also at 24 hours if the patient has received heparin (unfractionated or LMWH) in the previous 100 days). Patients with

platelet counts $<100 \times 10^9/l$ need to be monitored closely and if the platelet count falls below $50 \times 10^9/l$, thromboprophylaxis should be paused, except in very high-risk cases which should be discussed with a haemostasis expert.

The duration of thromboprophylaxis remains contentious. Clearly the risk of VTE falls as disease burden decreases, during which time there may be alterations in treatment (e.g. reduction in high dose dexamethasone dosing schedules). The majority of VTE in myeloma patients occur within the first 6 months of treatment. Thus, for example, thromboprophylaxis may be given for at least the first 4 to 6 months of treatment until disease control of active myeloma is achieved with more intensive regimens, and may then be de-escalated or discontinued. However, it is important that the strategy is individualized according to the presence of risk factors in each patient.

Table I Risk assessment model for the prevention of venous thromboembolism in multiple myeloma patients treated with thalidomide or lenalidomide (adapted from (Palumbo *et al*, 2008c)

Individual/ Myeloma risk factors	
Personal or family history of VTE	If no risk factors (RF) or only 1 RF consider aspirin If 2 or more RF present consider either: LMWH (high risk prophylactic dose eg. enoxaparin 40mg od) or Warfarin (target INR 2.5)
Obesity (BMI ≥ 30)	
Co-morbidities: cardiac, diabetes, renal impairment, chronic inflammatory disease	
Immobility (acute or chronic)	
Thrombophilias, myeloproliferative disorders, haemoglobinopathies	
Recent surgery (within 6 weeks): neuro-, trauma, orthopaedic, general, other	
Medications: erythropoiesis stimulating agents, HRT, tamoxifen/stilboestrol	
New diagnosis Myeloma	
Hyperviscosity	
Myeloma Therapy	
Doxorubicin	LMWH (high risk prophylactic dose eg. enoxaparin 40mg od) or Warfarin (target INR 2.5)
High-dose steroid ($\geq 480\text{mg/month}$ dexamethasone or equivalent)	
Combination chemotherapy	

Bleeding Risk Factors: the presence of a bleeding risk factor should prompt clinicians to consider whether bleeding risk is sufficient to preclude pharmacological thromboprophylaxis.
Active bleeding
Haemophilia or other known bleeding disorder
Platelet count $<100 \times 10^9/l$
Acute stroke in previous month (haemorrhagic or ischaemic)
Blood pressure $>200\text{mmHg}$ systolic or $>120\text{mmHg}$ diastolic
Severe liver disease (abnormal PT or known varices)

Severe renal disease (CrCl <30ml/min)
Undergoing procedure or intervention with high bleeding risk

3.2.3 Treatment of VTE in myeloma patients

As with other patients with suspected VTE, objective diagnosis should be made using appropriate imaging investigations, and treatment should follow best practice guidelines such as those of the American College of Chest Physicians (Kearon *et al*, 2008). No clear guidance can be given for the duration of anti-coagulation in myeloma patients developing VTE, but ongoing risk assessment is reasonable. There is good evidence that, in cancer patients, where the risk of VTE recurrence after discontinuation of anticoagulation is as high as 10%, long term maintenance with LMWH is significantly less frequently associated with recurrent VTE than warfarin, without an increased risk of bleeding (Lee *et al*, 2003; López *et al*, 2004). Based on these studies, extended therapy with LMWH should be considered in myeloma patients balancing these advantages against the inconvenience of daily injections, costs and long term complications. Those starting LMWH should have a baseline platelet count checked, and consideration given to checking further platelet counts every 2 to 4 days for the first 2 weeks of treatment, to screen for heparin-induced thrombocytopenia (HIT) (with a platelet count also at 24 hours if the patient has received heparin (unfractionated or LMWH) in the previous 100 days). For patients at extremes of body weight, or with renal impairment (Cr Cl <30mL/min) treatment dose LMWH may need to be dose adjusted, given in divided doses and monitored with anti-Xa levels.

If patients have a confirmed VTE whilst taking thalidomide or lenalidomide, it is reasonable to transiently discontinue these drugs until a fully anticoagulated state is established. Anticoagulation should be continued for the total duration of the treatment and then repeat VTE risk assessment performed to inform the duration of ongoing anticoagulation thereafter.

Recommendations

- **Cancer, cancer therapies, infection, previous VTE, immobility, obesity, paraplegia, ESA treatment, dehydration and renal failure are all well-recognised risk factors for venous thromboembolism (VTE), particularly in hospitalised patients. As**

with other areas of thromboprophylaxis, a risk stratified approach is appropriate in patients with myeloma (Grade C recommendation; level IV evidence).

- All patients who are due to start thalidomide or lenalidomide-containing therapy should undergo a risk assessment for VTE and prospectively receive appropriate thromboprophylactic measures. In patients receiving thalidomide or lenalidomide (Grade C recommendation; level IV evidence):
 - If no other VTE risk factors are present, aspirin 75-150 mg o.d. may be considered as VTE prophylaxis unless contraindicated (Grade C recommendation; level IV evidence).
 - If one or more major risk factors for VTE are present, prophylaxis with LMWH or adjusted therapeutic-dose warfarin is appropriate. This should be the consideration in most patients with active myeloma undergoing treatment with combination chemotherapy, unless contraindicated (Grade C recommendation; level IV evidence).
 - Patients with previous VTE should be considered for prophylaxis with adjusted therapeutic-dose warfarin or LMWH (high risk prophylactic dose should be considered). There is no role for fixed, low dose warfarin (Grade C recommendation; level IV evidence).
 - The duration of thromboprophylaxis remains unclear but guided by risk factors such as active disease (e.g. for the first 4 to 6 months of treatment until disease control achieved) and de-escalated or discontinued unless there are ongoing significant risk factors (Grade C recommendation; level IV evidence).
 - Treatment of confirmed VTE should follow current practice guidelines using adjusted dose warfarin or LMWH and appropriate monitoring (Grade C recommendation; level IV evidence).

4. Infection

Myeloma is associated with an increased incidence of early infection. This is related to deficits in both humoral and cellular immunity, reduced mobility and performance status which are all associated with both the disease and its treatment. It has been reported that up to 10% of patients die of infective causes within 60 days of diagnosis (Augustson *et al*, 2005). Neutropenia is not usually a factor in early infection (Augustson *et al*, 2005)

There is increasing evidence showing that high dose steroids in the elderly or in patients with poor performance may be detrimental with increased toxicity and a higher mortality rate in the short-term and consideration should be given to the use of lower doses in this group (Ludwig *et al*, 2009; Morgan *et al*, 2009; Rajkumar *et al*, 2009a). Patient education as well as access to 24 hour specialist advice and treatment is crucial in preventing and managing infection in myeloma.

Streptococcus pneumoniae, *Haemophilus influenzae* and Gram negative bacilli are the most frequent causes of infection in myeloma patients (Savage *et al*, 1982). Prophylactic antibiotics may have a role in reducing infection rates but have the potential for leading to increased *Clostridium difficile* infection and antibiotic resistance. The role of prophylactic antibiotics in myeloma needs to be addressed in larger numbers of multiple myeloma patients and their routine use is not recommended. Prophylactic immunoglobulin replacement has been shown to have some benefit in reducing infection rates in patients in plateau phase (Chapel *et al*, 1994) but no effect has been demonstrated in newly diagnosed patients (Salmon *et al*, 1967). A reasonable regimen is a dose of intravenous immunoglobulin of 500mg/kg administered every months for up to 6 months.

Studies in myeloma indicate suboptimal antibody responses to a variety of vaccines particularly worse for polysaccharide rather than protein antigens (Robertson *et al*, 2000). The value of influenza vaccination is unclear in myeloma although there is some evidence for efficacy in solid tumour patients undergoing chemotherapy (Melcher *et al*, 2005). Granulocyte colony stimulating factor (G-CSF) may have a role in reducing treatment-associated neutropenia (Mateos *et al*, 2008a) and following autologous peripheral blood stem cell transplantation (Sung *et al*, 2007).

Patient education as well as access to 24 hour specialist advice and treatment is crucial in preventing and managing infection in myeloma.

Recommendations

- **Vaccination against influenza, *Streptococcus pneumonia* and *Haemophilus influenzae* is recommended but efficacy is not guaranteed (Grade C recommendation; level IV evidence)**

- **Prophylactic immunoglobulin is not routinely recommended but may be useful in a small sub-set of patients with severe, recurrent bacterial infections and hypogammaglobulinaemia (Grade C recommendation; level IV evidence)**
- **Prophylactic aciclovir is recommended for patients receiving bortezomib therapy, following ASCT or patients with recurrent herpetic infections (Grade C recommendation; level IV evidence)**

5. Pain management

5.1 Prevalence and impact of pain in myeloma

Pain is one of the commonest symptoms experienced by myeloma patients and indeed it may have been the reason for initial presentation, or of subsequent relapse. Up to 67% of patients report pain at diagnosis, although this may have been present for several months before (Kariyawasan *et al*, 2007). At diagnosis, pain may be due to the disease process itself (predominantly from destructive bone disease, but occasionally from plasmacytomas directly affecting neural tissues), or it may signify a co-morbidity (eg degenerative arthritis or osteoporosis). Later in the course of the disease, pain often arises as a side-effect of therapies, eg thalidomide or bortezomib neuropathy.

Bone pain is associated with significant morbidity and impact on activities of daily living, especially if the spine or lower limbs are involved, impeding mobility. It is important to ask about these issues, as referral to OT or physiotherapy may be helpful alongside disease-directed therapy. Bone marrow examinations are usually associated with short-lived pain, but in some individuals this is sustained, perhaps because of neuroma formation in the biopsy site.

During intensive treatment, several pain syndromes have been identified (Niscola *et al*, 2006): these include deep somatic pain caused by growth factors and chemotherapy induced oropharyngeal mucositis.

Particularly in older patients, it is important to always consider co-morbidities such as arthritis or osteoporosis mimicking bony malignant pain; diabetes or carpal tunnel syndrome mimicking peripheral neuropathy; and post-herpetic neuralgia as a common cause of persistent pain.

Recommendation

- **All staff working with myeloma patients should be aware of the high incidence of pain and other symptoms, and be able to diagnose their cause (grade C recommendation, level IV evidence).**

5.2 Assessment of pain

It is important to make an accurate clinical assessment of pain, which starts with taking a history but may involve imaging by Xray, bone scan, CT or MRI. Patients can regularly be asked to score their pain on a 0-10 numerical scale (where 10 is the worst pain imaginable) at initial diagnosis and follow-up visits (Shi *et al*, 2009a). Patients unable to give a numerical score from 0-10 may find it easier to rate pain verbally on a four-point scale from 'none' to 'severe'. A numerical pain score of 5 or over is equivalent to moderate-severe pain and patients repeatedly scoring at this level should be referred to a palliative care or pain team (Serlin *et al*, 1995). Similarly, patients who do not achieve a reduction of 2 or greater on a 0-10 pain scale may benefit from specialist referral.

If the patient has more than one site of pain, these should all be assessed and recorded separately, as they may have different aetiologies and may require different interventions. A body chart is very helpful to record the sites of all pains, including radiation and dermatomal distribution for nerve root pain or the extent of stocking and glove distribution for peripheral neuropathy. Both the body chart and 0-10 scales for patient recording of pain at rest, on movement and the impact of pain on other activities and sleep are all captured by the Brief Pain Inventory – Short Form (Shi *et al*, 2009a).

To diagnose the presence of neuropathic pain, the LANSS scale can be used (Bennett *et al*, 2007). Specialist neurological consultation and electrophysiological testing are not usually required, unless there are unexplained dermatomal distributions.

Recommendations

- **Pain should be assessed regularly in myeloma patients at all stages of the disease (Grade C recommendation; level IV evidence)**

- **Pain should be measured using a 0-10 scale; alternatively it can be measured on a verbal none-mild-moderate-severe scale (Grade B recommendation; level III evidence)**
- **A reduction of 2 or greater on the 0-10 scale is clinically significant; if this is not achieved then the patient should be referred to a specialist (Grade C recommendation; level IV evidence)**
- **A body chart should be used to document sites of different pains (Grade B recommendation; level III evidence)**
- **The Brief Pain Inventory – Short Form should be the standard for full pain assessments (Grade A recommendation; level Ib evidence)**
- **The presence of neuropathic pain should be assessed using the LANSS scale (Grade A recommendation; level Ib evidence)**

5.3 Interventions

5.3.1 Introduction

In 1986 the WHO published its cancer pain programme which featured the 3-step analgesic ladder which has been widely used (Ventafridda *et al*, 1985). In recent years the limitations of the simplified WHO approach, in contrast to the advancements in pain science has been questioned (Ahmedzai and Boland 2007). Our understanding of cancer pain mechanisms and modes of action of analgesic drugs has increased greatly in recent years and a modern evidence-based approach based on the pathophysiological cause of specific types of pain is desirable (http://www.britishpainsociety.org/book_cancer_pain.pdf) (Raphael *et al*, 2010a; Raphael *et al*, 2010b). Such an approach would incorporate, for example, the molecular mechanism of the drugs which target neuronal or synaptic transmission and awareness of the toxicities of drugs (Ballantyne and Mao 2003; Holdcroft and Power 2003; Mantyh *et al*, 2002). Cancer pain usually requires a combination therapy approach which can include an opioid, a calcium channel blocker, sodium channel blockers and noradrenaline reuptake inhibitors (SNRIs or tricyclic antidepressants) (Ossipov and Porreca 2005). Depending on the experience of the haematologist and/or general practitioner, optimal treatment with systemic calcium and systemic sodium channel

blockers may require prescription by or advice from a pain or palliative medicine specialist. NSAIDs have limited use in myeloma because of the potentially serious renal toxicity.

In addition, modern pain management for myeloma should offer a range of non-pharmacological interventions including bisphosphonates, radiotherapy, percutaneous vertebroplasty and balloon kyphoplasty, orthopaedic fixation of spine and long bones (Siemionow and Lieberman 2008). The latter surgical interventions should be considered early in the course of disease in order to stabilise long bone and the vertebral spine from fracture and the consequent adverse effects on mobility and quality of life. For selected patients with lack of response or intolerable toxicity from systemic opioids, spinal delivery – preferably via an implanted intrathecal catheter – should be considered (Smith *et al*, 2005). Some patients, particularly those with high levels of anxiety, also benefit from psychological techniques such as relaxation therapy.

Recommendation

- **Pain arising in myeloma patients should be managed using a multi-modal, mechanism-based approach including evidence-based pharmacological therapies alongside non-drug methods such as radiotherapy, bisphosphonates, and where appropriate, interventional and psychological techniques (Grade B recommendation; level III evidence).**

5.3.2 Non-opioid analgesics

Paracetamol is a useful analgesic in cancer-related pain and other chronic pains and should be prescribed at a dose of up to 1 gram qid (p.o. or i.v. in patients who cannot take oral medication, e.g. because of vomiting or mucositis).

NSAIDs should be avoided apart from very short term use (eg 3-5 days) with acute severe pain, eg bone fracture. They should not be used in the presence of renal impairment, and used with extreme caution in myeloma patients in view of the risk of precipitating renal compromise.

Recommendations

- **Paracetamol can be prescribed up to 1g qid (Grade B recommendation; level III evidence)**

- **NSAIDs should generally be avoided in myeloma. (Grade C recommendation; level IV evidence)**

5.3.2 Opioid analgesics

5.3.2.1 Mild pain (0-4 on a 0-10 scale)

For patients with mild pain (<5/10), and those who are opioid-naïve, normal release tramadol is a reasonable choice of analgesic agent. Tramadol has 1/5th the potency of oral morphine and the starting dose is 50mg 6 hourly prn or qid. Patients who respond but need qid dosing are best placed onto the bd sustained release form. Codeine can also be used but it is a pro-drug of morphine, and 10-15% of the population is unable to convert it into active morphine, leaving them with unacceptable toxicity (Lötsch and Geisslinger 2006).

5.3.2.2 Moderate to severe chronic pain (>4/10)

Patients with chronic moderate (5-6/10) or severe pain (>6/10) can be started on tramadol as above, but will usually need to go onto more potent opioids rapidly if they do not respond. Oxycodone is twice the potency of morphine and is associated with less drowsiness and hallucinations. For rapid onset, the normal release preparation can be used 4-6hourly or qid, but most patients eventually prefer the convenience of the bd sustained release forms (Mucci-LoRusso *et al*, 1998).

Patches can be used to deliver either fentanyl or buprenorphine, both of which are very potent opioids. Fentanyl causes significantly less nausea, sedation and constipation compared to morphine (Clark *et al*, 2004). When given the choice of fentanyl patches or oral morphine for chronic pain, patients prefer the patches (Ahmedzai and Brooks 1997). Buprenorphine often initially causes nausea but this can be covered by the use of an anti-emetic such as metoclopramide and is otherwise well tolerated. Note that there are two formulations of buprenorphine patch – the low dose (5-10mcg/h) are primarily for use in arthritis or other non-cancer patients, whereas the higher doses (35mcg/hr and greater) are more useful for cancer pain.

When using normal release oral medication, the dose can be titrated up daily by 30-50% until pain is controlled or unacceptable side effects occur. With sustained release oral medication it is advisable to wait 2-3 days between dose increments. With patches, doses should not normally be increased at less than 3 day intervals.

With all sustained release analgesics, it is essential to offer the patient a normal release 'rescue medication' for breakthrough pain. This is particularly important when breakthrough pain occurs quickly and predictably, e.g. on weight-bearing with disease in the spine or legs. It is important to distinguish this kind of 'incident pain' from pain arising from end of dose failure with sustained release medications, or spontaneous pains associated with neuropathy or opioid-induced hyperalgesia (Davies *et al*, 2009). Normal release oxycodone or morphine can be used, at 1/6th of the current 24 hour total opioid dose. However, often the absorption of these oral drugs can be too slow for some episodes of breakthrough pain. Fentanyl has a high bioavailability via the transmucosal route, which has led to the development of fast-acting (but short-lived) fentanyl formulations. These include fentanyl lozenges (Actiq); buccal tablets (Effentora); or sublingual tablets (Lennernäs *et al*, 2010; Weinstein *et al*, 2009). Nasal sprays will also soon be available. Normally, a patient should not need to use more 2-3 of these relatively expensive fentanyl formulations per day for breakthrough pain; if more are being taken, either the background medication needs to be increased or the patient should be referred to a specialist.

There is no place for pethidine in the treatment of pain in myeloma.

5.3.2.3 Acute onset moderate-severe pain (>4/10)

For patients with sudden onset of moderate to severe pain, e.g. after long bone fracture or vertebral collapse, the subcutaneous route is recommended, e.g. with oxycodone or morphine, until the pain is controlled and then a sustained release oral preparation can be started. Opioids can be combined with other drugs, e.g. an anti-emetic in the syringe driver. There is usually no advantage of using the intravenous route, except in acute severe pain and this should be administered by a pain or palliative medicine specialist. Patient controlled analgesia (PCA) pumps are also of little value in myeloma patients, except in some cases of very severe oropharyngeal mucositis.

For non-emergency situations, there is no advantage of using injections or subcutaneous syringe drivers, except when the patient is vomiting or otherwise unable to take oral medication.

Diamorphine is a pro-drug for morphine and apart from its greater water solubility, it has no advantages over morphine or oxycodone for injections.

5.3.2.4 Adverse effects of opioids

With all opioids, it is important to offer the patient a laxative and to keep checking for the development of constipation. Transdermal fentanyl and buprenorphine are associated with reduced incidence of constipation (Clark *et al*, 2004; Tassinari *et al*, 2009). It is not necessary to routinely prescribe an anti-emetic with opioids, except for the first week when starting buprenorphine.

Most opioids cause dose-related sedation; however, fentanyl and oxycodone are associated with reduced sedation compared to morphine (Ahmedzai and Brooks 1997; Clark *et al*, 2004; Reid *et al*, 2006). Patients who experience intolerable sedation due to opioids (or other drugs, eg thalidomide) may be considered for a trial of a psychostimulant such as methylphenidate or modafanil; this should only be prescribed by a specialist in palliative medicine. (See section on Fatigue in section 8 'Other symptoms'.)

Respiratory depression is uncommon in patients treated chronically with opioids as long as dose increments are made carefully as outlined above. With the initiation of opioids, it is common to see a reduction in respiratory rate; however, this is usually balanced by changes in tidal volume so that minute ventilation initially remains steady. Care needs to be taken in patients with COPD or obstructive sleep apnoea, in whom the respiratory depression can occur even with low doses of opioids. True respiratory depression caused by opioids is diagnosed by a reduction in oxygen saturation ($\text{SaO}_2 < 90\%$) or by arterial blood gases. If this occurs, naloxone can be given but care must be taken not to provoke a serious increase in pain. Advice on future opioid dosing should be sought from a specialist in pain or palliative medicine.

Recently a condition known as opioid-induced hyperalgesia has been consistently identified in animal studies and has also been demonstrated to occur in human studies (Ballantyne and Mao 2003). This condition is characterised by increasing reporting of pain

in the presence of increasing opioid dosage. The pain can be localised to the original lesion but is often generalised to adjacent dermatomes. The skin in the affected area may show hyperalgesia (increased pain response on normal painful stimulus) or allodynia (pain felt even on light touch). This is thought to be caused by downstream intracellular signalling mechanisms from the activated opioid receptor and involves the induction of nitric oxide and the opening of NMDA channels, which are responsible for maintaining chronic and neuropathic pain (Mao 2008; Mao *et al*, 2002). The management of this condition should be left to specialists in pain or palliative medicine; it involves reduction in the opioid dosage along with the introduction of an NMDA channel blocker such as ketamine or methadone.

For details on currently available opioids and their usual starting doses; dose equivalents for conversions; calculation of prn dosing , see (Twycross and Wilcock 2008).

Recommendations

- **For mild-moderate pain (<5/10), oral tramadol is a recommended (alternative is codeine) (Grade C recommendation; level IV evidence)**
- **For chronic moderate-severe pain (>4/10), oxycodone is recommended, either normal or sustained release (alternative is morphine) (Grade B recommendation; level III evidence)**
- **For patients with chronic moderate-severe pain, fentanyl or buprenorphine patches are convenient, cause fewer side-effects and are preferred by patients (Grade A recommendation; level Ib evidence)**
- **For acute severe pain (>6/10) it is reasonable to initiate subcutaneous opioid therapy to achieve quicker pain control, using oxycodone injection (morphine is alternative) (Grade C recommendation; level IV evidence)**
- **For non-oral medication, eg patient is vomiting, has reduced consciousness or is moribund, the subcutaneous route is recommended (Grade C recommendation; level IV evidence)**
- **Patients on opioids should be regularly screened for toxicity - the main treatable adverse effects being constipation, emesis and sedation (Grade C recommendation; level IV evidence)**

- **All patients on opioids should be prescribed laxatives (Grade A recommendation; level I? evidence)**
- **Patients who develop drug-related sedation should be referred to a specialist for consideration of a psychostimulant (Grade B recommendation; level III evidence)**
- **Patients on opioids who experience increasing pain in spite of increasing doses and in the absence of increasing disease, should be considered as having opioid tolerance or opioid-induced hyperalgesia and should be referred to a pain or palliative medicine specialist (Grade C recommendation; level IV evidence)**

Table II Key points to compare opioids for analgesia used in UK

Opioid	Receptors	Routes	Equivalent 24 hour dose (oral morphine as reference point)	Advantages	Disadvantages	Renal failure	Hepatic failure	Other information
Morphine	MOR agonist	PO, IV, SC, Sp	30mg PO = 15mg SC, IV	Familiarity; Cheap; Available as normal release and modified release tablets	Sedation; Hallucinations; Nausea; Constipation; Immunosuppression	Do not use - high risk of toxic metabolites in renal failure	Caution if prothrombin time prolonged	Nausea usually side-effects are usually persistent
Tramadol	MOR agonist; SSRI	PO	150mg PO (maximum recommended dose is 400mg per 24 hours)	Less constipation than morphine; SNRI action may help with neuropathic pain	Only available PO May be immune-enhancing	Caution	Caution	Efficacy highly dependent on CYP2D6 phenotype; SNRI activity; Risk of serotonin syndrome
Codeine	MOR agonist	PO	300mg PO (NB – this is higher than maximum daily dose of 240mg)	Familiarity; OTC	Constipation; Sedation	Unsafe - as for morphine	Caution - as for morphine	Pro-drug of morphine; dependent; Often combined with paracetamol
Oxycodone	MOR, KOR agonist	PO, IV, SC	15mg PO = 7.5mg SC, IV	Reduced sedation, hallucinations <i>cf</i> morphine	Oral liquid unpleasant taste – use capsules; For parenteral use, concentrate (50mg/ml) now available as substitute for diamorphine); Probably immune-neutral	Caution – plasma concentrations may rise	Safe	Females have higher CYP3A4 and dependent
Fentanyl	MOR agonist	TM (buccal, sublingual, nasal), TD, IV, Sp	12mcg/h TD patch over 72 hours	Reduced sedation, emesis, constipation <i>cf</i> morphine; Convenience of 3 day patch	TM applications very short-acting (1-2 hours) – best reserved for incident (movement-related) pains and dressing changes, etc; Probably immunoneutral	Safe in renal failure (Alternative for parenteral use is alfentanil)	Safe	Use rapid action with caution in patients with addictive tendencies; Affected CYP3A4
Buprenorphine	Partial MOR, ORL1 agonist; KOR, DOR antagonist	TM (sublingual), TD, IV	20mcg/h TD patch over 7 days	Reduced respiratory depression; Convenience of patches (5-20mcg dose TD patches can last 7 days)	Nausea with initiation of higher dose TD patch; TM tablet causes nausea	Safe	Safe	Ceiling dose for respiratory depression may be lower in COPD patients; reverse other effects with naloxone; therapeutic dose
Diamorphine	MOR agonist	IV, SC	10mg SC, IV	Familiarity; High water solubility	As for morphine	Unsafe - as for morphine	Caution -as for morphine	Pro-drug of morphine; Has no advantage; high water solubility; that oxycodone is available in UK

Please note that this table has been compiled by one of the authors (SHA) as a guide for haematologists on how to use the common opioids available in the UK. Other opioids eg methadone and hydromorphone, are for specialist use by palliative medicine and pain management services. The new fast-acting transmucosal fentanyl products are also usually restricted to specialist initiation only. If in any doubt, eg possible interaction, renal failure, use of syringe drivers etc, and for further help in using

opioids with myeloma patients, the haematologist should contact these local specialist services or pharmacy.

Key

MOR, mu-opioid receptor; KOR, kappa-opioid receptor; DOR delta-opioid receptor; ORL1, opioid-receptor-like receptor; SSRI, selective serotonin reuptake inhibitor

PO, oral; IV, intravenous; SC, subcutaneous; Sp, spinal (epidural or intrathecal); TD, transdermal; TM, transmucosal (sublingual, buccal or nasal)

Note that although there is a perception that there is no effective ceiling dose for morphine or other strong opioids, it is advised to seek specialist advice when *total daily doses* reach these levels – morphine 200mg PO, 100mg parenteral; oxycodone 100mg PO, 50mg parenteral; buprenorphine patch 70mcg/hr; fentanyl patch 50mcg/hr; diamorphine 50mg parenteral

Main sources: ref Twycross 2008; ref Raphael et al 2010a; British National Formulary <http://bnf.org/bnf/index.htm> (last accessed 17 September 2010)

5.3.4 Special pain situations (see section 6 for neuropathic pain)

With all chronic cancer-related pains and iatrogenic neuropathies, there is a good case for using a calcium-channel blocking agent, eg gabapentin or pregabalin. However, gabapentin is associated with a risk of bone marrow suppression and should be used with caution during stem cell transplantation procedures (there is presently no evidence regarding pregabalin). Both gabapentin and pregabalin should be used at a reduced dose in renal failure.

For procedural pain, e.g. patients with bone fractures who are unable to lie down for MRI scanning or radiotherapy, it can be helpful to refer to the local specialist palliative care or pain service, who may use drugs such as ketamine or propofol.

For patients with continuing severe (>6/10) pain or those who are unable to tolerate analgesics because of adverse effects, help should be sought from a specialist service such as the palliative care team or chronic pain team. They will advise on and supervise use of options such as ketamine, methadone or spinal analgesia.

Recommendation

- **All patients with chronic pain should be considered for a calcium channel blocker (gabapentin or pregabalin) and a sodium channel blocker (lidocaine,**

oxcarbazepine) and a SNRI (duloxetine or amitriptyline). (Grade B recommendation; level III evidence)

5.4 Collaboration with specialist services

5.4.1 Palliative Medicine and Pain Services

Haematology teams should readily seek to share care of pain and other symptoms with local palliative and supportive care teams and radiotherapy services. Patients at home can be seen by community or hospice-based palliative care teams. Hospital chronic pain teams should be consulted for severe pain if palliative and supportive care teams are not available. Acute pain teams may be helpful if the patient has an acute severe pain, eg bone fracture causing immobilisation, which may respond to interventional procedures, e.g. local nerve blockade or spinal delivery of opioids and local anaesthetic. Orthopaedic surgeons or interventional radiologists are able to perform cement vertebroplasty or kyphoplasty for uncontrolled pain arising from vertebral collapse (see section 4.4.3). Psychologists can help with patients who have severe anxiety overlying pain and with other issues, eg phobias. Specialist nurses in haematology teams may have some training in symptom palliation but they should work and consult with palliative care specialist nurses and teams.

Warning signs indicating need to refer to local specialist service include:

- a. Increasing pain in spite of following above guidelines (especially if opioid tolerance or opioid-induced hyperalgesia is suspected)
- b. Intolerable adverse effects of standard treatments
- c. Additional psychological stresses or existential problems (eg coping with news of recurrence or facing death) which are preventing patients from adapting to or coping with symptoms
- d. Social or family circumstances that prevent the patient from adhering to the treatment regimen or being discharged from acute care

The median dose of morphine (or equivalent median dose of other opioids) in patients with advanced cancer who are being treated by specialists in palliative medicine is

approximately 120mg per day. It is therefore recommended that if the total daily dose of opioid in a myeloma patient exceeds this median dose, this should be a trigger for the patient should be discussed with, or referred to, the local palliative care team. (This cut-off dose is equivalent to approximately 60mg bd of oral morphine; 30mg bd of oral oxycodone; 25mcg/h of fentanyl patch). However, it should be noted that in many centres, the palliative care team may be too small to cope with a large number of such referrals; or patients who are still having potentially 'curative' or 'life-prolonging' treatments may fall outside their remit.

It is important when referring to a specialist palliative care service for symptom control to reassure the patient that this does not mean that they are at 'the end of life', since that is often the assumption of the public. The best way of supporting patients is shared care between the haematology department and the supportive and palliative care team or pain service.

5.4.2 Radiotherapy for pain relief

Local radiotherapy is effective for pain relief for skeletal disease and may also palliate soft tissue disease. Relatively low-radiation doses are required with a correspondingly low incidence of side effects and with the potential for re-irradiation for local recurrence of symptoms if required. Published dose–response data specific to myeloma are very limited. Mill and Griffith reported on 278 radiation fields in 128 patients treated over a wide dose range (Mill and Griffith 1980). Pain relief occurred in 91% (21% complete) at a median dose of 10–15 Gy in 2–3 Gy fractions. Only 6% of sites required re-treatment and this was unrelated to initial dose. Leigh analysed 316 sites in 101 patients; 97% experienced pain relief (complete in 26%) with a mean dose of 25 Gy (range 3–60 Gy); 6% relapsed after a median interval of 16 months and were retreated (Leigh *et al*, 1993). There was no correlation between dose and either response or relapse.

Myeloma is at least as radio-responsive as other types of skeletal metastases and it is reasonable to extrapolate guidance on radiotherapy dose and fractionation for pain relief in myeloma from randomised trials of localised radiotherapy for the treatment of bone metastases from a variety of primaries. A meta-analysis of 16 such trials concluded that there was no significant difference in complete or overall pain relief between single and multi-fraction regimes and no evidence of a dose response relationship between 8 Gy

single exposure and up to 40 Gy fractionated, although more frequent re-irradiation was noted in the lower dose arms (Wu *et al*, 1993).

Patients should have sufficient analgesia to cover the administration of radiotherapy, including the positioning necessary for its delivery and for the period following treatment until maximum benefit is achieved. In some cases it may be necessary to call on the services of the local pain or palliative care service to give acute analgesia such as ketamine to enable patients to lie on the table for treatment. Given that some but not all elements of pain in myeloma are radio-responsive an early reappraisal of symptomatic control measures is important.

Recommendations

- **Local radiotherapy is helpful for pain control; a dose of 8 Gy single fraction is recommended (Grade C recommendation; level IV evidence).**

5.4.3 Specialised spinal services

5.4.3.1 Introduction

Although the management of spinal pain is often conservative, in the absence of instability/neurological compromise, spinal or interventional radiological advice should be sought in cases of persistent/refractory pain. Vertebroplasty and kyphoplasty are alternative options for controlling pain associated with vertebral collapse. Vertebroplasty and kyphoplasty are both vertebral body augmentation techniques of percutaneous injection of bone cement to the vertebral bodies. They are best performed soon after the vertebra collapses and may be ineffective if many months have elapsed. Both techniques carry the small risk of cement leakage leading to pulmonary embolism and neural compromise. It is therefore important that there is access to a spinal surgery service when these procedures are performed.

5.4.3.2 Vertebroplasty

This involves the percutaneous injection, under general anaesthetic and i.v. sedation, and using radiological imaging, of polymethacrylate bone cement or equivalent biomaterial into

the vertebral body (Jensen and Kallmes 2002). Several vertebrae can be treated simultaneously. The injection allows local pain relief and bone strengthening but will not restore vertebral height. No randomized studies on the use of vertebroplasty in myeloma have been published. However, a recent review of 67 cases demonstrated improvements in pain (89%), mobility (70%) and use of narcotic analgesia (65%) (McDonald *et al*, 2008).

5.4.3.3 Kyphoplasty

This involves the percutaneous insertion of a small, inflatable balloon into the vertebral body; when inflated it produces a potential space. The balloon is then removed and bone cement is injected to fill the cavity. Although more time consuming than vertebroplasty the complication rates appear lower with similar potential benefits of both pain relief and improved function to vertebroplasty but with reduced risk of cement leak. There is also the potential to restore vertebral height but this only occurs in a minority of patients. At the present time, the documented use of kyphoplasty in myeloma is limited to case reports and small case series (Fourney *et al*, 2003; Masala *et al*, 2004) although outcomes in myeloma do appear comparable to those in osteoporosis (Lane *et al*, 2004). The UK National Institute for Clinical Excellence (NICE) (2003a) reviewed the evidence on kyphoplasty for vertebral compression fractures and made the following recommendations:

- The procedure should only be undertaken where there are good arrangements for access to spinal surgery services in the event of complications.
- The indication and suitability for the procedure should be agreed by the multidisciplinary specialist team.
- Clinicians performing the procedure must have received training to an appropriate level of expertise.
- The procedure should be limited to patients whose pain is refractory to more conservative treatment.

Recommendations

- **The use of vertebroplasty or kyphoplasty may be considered in patients with persistent pain (Grade B recommendation; level III evidence).**
- **The use of kyphoplasty should follow the NICE recommendations summarized above (Grade C recommendation; level IV evidence).**

6. Management of peripheral neuropathy

6.1 Scope

Many patients with myeloma have subclinical or even clinical peripheral neuropathy at diagnosis, often due to co-morbidities (see 5.2.2). These patients are at risk of worsening peripheral neuropathy when exposed to potentially neurotoxic drug treatments, such as thalidomide and bortezomib. The cause of PN in myeloma patients is multifactorial and when patients are assessed, it is important to grade the degree of neuropathy using a recognised scale such as the National Cancer Institute (NCI) Common Toxicity Criteria (Trotti *et al*, 2003), LANSS (Bennett 2001) or the Total Neuropathy Score (Cavaletti *et al*, 2007). Additionally, it may be helpful to evaluate the pain intensity and its effect on functioning, using the 'Worst pain in past week' item (score 0-10) of the Brief Pain Inventory (Shi *et al*, 2009b).

6.2 Causes

Peripheral neuropathy in myeloma patients can be subdivided as follows:

6.2.1 Disease or M protein associated peripheral neuropathy

Spinal cord or nerve root compression is a common neurological complication of myeloma due to compression by plasmacytoma, lytic or extramedullary disease (Silberman and Lonial 2008) and requires appropriate imaging and specific treatment including a specialist opinion as to the need for surgical intervention or radiotherapy.

The well recognised association between MGUS and PN (Kelly *et al*, 1981; Nobile-Orazio *et al*, 1992) is addressed in the Guidelines for Management of MGUS (Bird *et al*, 2009). The incidence of PN in myeloma is less well documented, with earlier studies indicating that symptomatic PN is present in 3 to 13% at diagnosis (Silverstein and Doniger 1963; Walsh 1971) and subclinical PN in one-third of patients, detectable by nerve conduction or histopathological studies (Kelly *et al* 1981). The reported prevalence of sensory PN may depend on the study cohort, the methods of detection and the criteria used, with a recent study reporting rates of pre-treatment sensory PN in up to 20% of patients, and neuropathic abnormalities in as many as 54%. (Richardson *et al* 2009). The cause of the neuropathy in many cases of myeloma is not clear and may be multifactorial, and studies

have also varied in relation to rates of small or large fibre or mixed PN. In those cases where amyloidosis and toxicity due to chemotherapy are not the cause, the M protein itself or other consequences of the underlying disease may play a part. Clinically, a symmetrical, distal sensory/motor neuropathy inducing paraesthesiae and numbness in the hands and feet is seen. Treatment of the myeloma results in varying degrees of improvement in the PN in this situation and in some cases the PN may worsen (Kyle 1992). Given the variation between studies, it is likely that further studies with standardised criteria are warranted to characterise baseline PN in patients with myeloma.

POEMS syndrome and AL amyloidosis are more specialised situations, PN is a significant clinical feature in 85 to 100% of patients affected by POEMS syndrome (Dispenzieri and Gertz 2004). It is a consequence of axonal degeneration and demyelination, typically distal, symmetrical and initially sensory, but as the condition progresses, a disabling symmetrical weakness may develop.

PN affects 17% of patients with AL amyloidosis at diagnosis (Kyle and Gertz 1995). The PN is typically axonal and characteristically painful, distal and symmetrical and often associated with an autonomic neuropathy. Treatment options for AL amyloidosis are discussed in the BCSH Guidelines (Guidelines Working Group of UK Myeloma Forum (2004a)). Cryoglobulinaemia is another recognised source of PN.

6.2.2 Peripheral neuropathy related to co-morbidities

Conditions such as diabetes mellitus, carpal tunnel and other nerve compression syndromes, including chronic inflammatory demyelinating polyradiculoneuropathy, chronic renal failure and vitamin B12 deficiency should be actively sought and appropriately managed, with specialist input as needed.

6.2.3 Chemotherapy induced peripheral neuropathy (CIPN)

Chemotherapy induced peripheral neuropathy, also known as treatment emergent peripheral neuropathy, is a major aspect of myeloma management. CIPN has been a long recognised complication of vinca alkaloid and platinum-based treatments (Wolf *et al*, 2008) and may be significantly dose limiting, but these drugs are no longer in regular use in myeloma. There is emerging evidence for the incidence and natural history of PN due

to novel therapies, including thalidomide-induced PN (TiPN) (Palumbo *et al*, 2008b) and bortezomib-induced PN (BiPN) (Argyriou *et al*, 2008), which may be considered as distinct clinical entities

TiPN may arise after prolonged administration of thalidomide (39–75% of patients of patients treated for 12 months), is mostly mild to moderate in severity (Mileshkin *et al*, 2006) and appears to be a cumulative effect (Cavaletti *et al*, 2004). Initial symptoms include sensory changes such as paraesthesia and hyperaesthesia, motor symptoms and autonomic dysfunction. Later effects include loss of vibration and joint position sense, which may lead to ataxia and progressive gait disturbance. Nerve conduction studies do not reliably predict the onset of significant TiPN and do not necessarily correlate with the clinical findings (Mileshkin *et al*, 2006). Reduction or temporary discontinuation of the drug usually leads to a clinical improvement in the symptoms whereas continuation of dose intense treatment in the face of neuropathy may cause permanent neurological damage. Mileshkin and other investigators have recommended that thalidomide therapy should not exceed 6 months as the risk of TiPN are unacceptably high.

BiPN is characterised by neuropathic pain and a length-dependent distal sensory neuropathy with suppression of reflexes (Cavaletti and Nobile-Orazio 2007). Motor neuropathy may follow and infrequently results in mild to severe distal weakness in the lower limbs (El-Cheikh *et al*, 2008). There may also be a significant autonomic component, which manifests as dizziness, hypotension, diarrhoea or constipation and/or extreme fatigue. It is thought to occur at a certain threshold (within 5 cycles but rarely beyond) of treatment (Richardson *et al*, 2006b) and may be more likely to occur within the setting of renal impairment in keeping with other therapy-related toxicities in this setting (Jagannath *et al*, 2005a). Electrophysiological testing reveals a mainly distal sensori-motor axonal loss, with secondary demyelination. The symptoms of BiPN improve or completely resolve in the majority of patients after a median of 3 months following discontinuation of the drug (Badros *et al*, 2007; Richardson *et al*, 2003; Richardson *et al*, 2006b), but in a proportion of cases, symptoms have taken up to 2 years to improve (El-Cheikh *et al*, 2008). Apart from a graded dose reduction or withdrawal (Richardson *et al*, 2006b), the only treatment for BiPN is symptomatic relief. No effective prophylactic treatment is available and any use of nutritional supplements should be restricted to low doses to avoid harm from excessive doses of pyridoxine (Levine and Saltzman 2004). In particular, caution should be exercised

with supplements containing ascorbic acid which may inhibit the anti-myeloma effect of bortezomib (Peronne et al 2009).

The frequency of BiPN varies between studies. Data from the SUMMIT and CREST phase II trials in the relapsed setting suggest that BiPN occurs in about 35% of patients (13% G3 and 0.4% G4), and led to discontinuation of therapy in 5% and dose reduction in 12% (Badros *et al*, 2007; Richardson *et al*, 2006b). Other studies suggest a higher incidence in up to 75% of patients with relapsed disease and 33% of newly diagnosed patients, while grade 3 to 4 neurotoxicity may affect 30% of relapsed and 18% of newly diagnosed patients who receive bortezomib (Badros *et al*, 2007; Jagannath *et al*, 2005b).

6.3 Assessment and investigation of peripheral neuropathy

An accurate neurological history should be taken from all patients prior to commencement of neurotoxic agents and regularly during the course of therapy (Table III). Patients should be reviewed in person at the start of each cycle to ensure that emergent symptoms are detected and acted upon. Dose-reductions may be needed within a treatment cycle if symptoms are progressive, so as to avoid the irreversible neurological damage that may result from waiting until the next cycle to make a change.

Table III. Important features of the neurological history

<u>Sensory Symptoms</u>	<u>Motor Symptoms</u>	<u>Autonomic Symptoms</u>
A sensation of wearing an invisible "glove" or "sock" Wooden quality Numbness, pins & needles Feeling of walking on pebbles Feeling of tightness & swelling Burning sensation or freezing pain Sharp, jabbing or electric shock-like pain Extreme sensitivity to touch Usually worse at night Loss of balance and coordination Cramps in the feet and calves	Tripping on the toes Loss of grip strength	Orthostatic dizziness Constipation Diarrhoea Bladder incontinence Sexual dysfunction Dry eyes Dry mouth

It is important to grade the severity of the symptoms of CIPN using a scale, such as the NCI-CTC (Trotti *et al*, 2003), to provide an objective assessment of the PN, so that if different members of the team are assessing the same patient, sequential assessments can be made and trends identified.

Initial investigations should be tailored according to the history and examination and may include those listed in Table IV. Vitamin B12 deficiency should be screened for periodically. Metabolic and autoimmune causes should also be considered. If there are prominent features of small fibre neuropathy, then AL amyloidosis should be excluded by tissue biopsy or serum amyloid P (SAP) scan; any further investigations such as electrophysiological studies or cerebrospinal fluid protein estimation should be directed by a neurologist.

Table IV. Investigation of peripheral neuropathy

Routine	Additional	Consider	Specialist tests

FBC U&E LFT Ca ThFT	Serum B12 Lipids Fasting glucose ANA ENA ANCA Serum ACE	MRI of spine and plexii (if radicular compression is suspected) Oral glucose tolerance test SAP scan	Nerve conduction studies Lumbar puncture and CSF protein Anti-neuronal antibodies Nerve biopsy
---------------------------------	---	--	--

Abbreviations: FBC: Full Blood Count; U&E: urea and electrolytes; LFT: liver function tests; Ca: Serum calcium; ThFT: thyroid function tests; ANA: anti nuclear antibody; ENA: extractable nuclear antigen; ANCA: Antineutrophil cytoplasmic antibody; ACE: angiotensin converting enzyme; MRI: magnetic resonance imaging; SAP: serum amyloid P; CSF: cerebrospinal fluid

6.4 Management of peripheral neuropathy

The management of peripheral neuropathy should include symptom control along with treatment of any potentially reversible causes. Identification and correction of Vitamin B12 deficiency is important and optimal management of co-morbid causes such as diabetes mellitus or alcohol excess may also improve tolerance of neurotoxic drugs. An awareness of the spectrum of symptoms that herald CIPN is crucial. Such symptoms need to be carefully sought at each meeting with the patient.

Careful monitoring of patients receiving bortezomib and prompt dose and schedule modifications are essential. Dose reductions according to the SPC, or changing from twice weekly to once weekly dosing should be instituted as soon as symptoms emerge. Temporary interruptions in therapy may also be beneficial, before resuming on a new schedule/dose. Recent data from front line protocols incorporating bortezomib suggest that a weekly regimen is as effective and associated with less neuropathy than twice weekly regimens (Mateos *et al*, 2008; Palumbo *et al*, 2008a, Brinthen *et al* 2010). Continuation of dose intense treatment in the face of neuropathy may cause permanent neurological damage. Measurement of lying and standing blood pressures weekly in patients receiving bortezomib, may detect autonomic neuropathy before it becomes a debilitating problem for the patient. The administration of intravenous normal saline prior to each dose of bortezomib may improve tolerance of the drug.

Neuropathic pain is often poorly responsive to standard analgesic regimes. There has been very little research specifically in the management of chemotherapy-induced painful peripheral neuropathy, and that has mostly been in solid tumours (Tsavaris *et al*, 2008). Opioids can be effective but if used alone in high dose are associated with significant adverse effects (Rowbotham *et al*, 2003). A multi-modal approach using opioids together with other pain modulating drugs is now recommended (Ossipov and Porreca 2005; Raphael *et al*, 2010b). Thus a calcium channel blocker should be added early (eg gabapentin or pregabalin); it may be necessary to add a sodium channel blocking agent, e.g. oxcarbazepine (carbamazepine should be avoided because of drug interactions); or an SNRI, e.g. amitriptyline or duloxetine (Jensen *et al*, 2009).

Several studies have shown that adding gabapentin to an opioid in patients with cancer-related neuropathic pain can give improved analgesia with reduced adverse effects compared to using either agent alone (Caraceni *et al*, 2004; Ho *et al*, 2009; Keskinbora *et al*, 2007). The study of chemotherapy-induced peripheral neuropathy from (Tsavaris *et al*, 2008) was not randomised but 75 patients who received 800mg /day of gabapentin were compared to 35 patients who declined gabapentin but received naproxen and codeine/paracetamol instead. Gabapentin led to a complete response in 25.3% of patients (19/75), partial response in 44% (33/75), minor response in 25.3% (19/75), and no response in 5.3% (4/75). In the “control group”, none experienced complete response (0/35), while partial, minor, and no response were observed in 5.7% (2/35), 45.7% (16/35), and 48.6% (17/35), respectively. The response to gabapentin correlated with the severity of the underlying neurotoxicity. Approximately 25% of patients receiving gabapentin experienced mild somnolence, but none discontinued it. Note that gabapentin may be associated with myelosuppression and so should be avoided around the time of stem cell transplant.

The haematologist who is not familiar with these agents should seek advice from the local chronic pain or palliative care service. For patients with continuing severe pain in spite of initiating these drugs or those who are unable to tolerate analgesics because of adverse effects, specialist help is essential. They will advise on dose modifications and can also initiate specialist options such as ketamine, methadone or spinal analgesia.

In addition, topical treatments may be of benefit. Capsaicin cream 0.075% acts on peripheral nerve TRPV1 heat and pain receptors; menthol acts on TRPM8 receptors for cold and may both be helpful in patients with ‘cold’ or ‘hot’ dysaesthesia respectively

(Vriens et al 2008). Emollients, such as cocoa butter, may help some patients but the physiological mechanism is unclear. In other forms of superficial neuropathic pains (e.g. post-herpetic neuralgia or scar pain), the sodium channel blocker lidocaine can be used topically as a 5% plaster, applied to the affected area for 12 hours and then left off for 12 hours. Some patients obtain relief within a few days but the peak effect is reached with 2-4 weeks (Binder et al 2009, Baron et al 2009).

Recommendations

- **Symptoms and signs of peripheral and autonomic neuropathy should be actively sought using a validated screening tool, such as LANSS, and the cause identified, where possible (Grade C recommendation; level IV evidence)**
- **Clinical evidence of a significant (e.g. >NCI grade 2) or progressive PN at diagnosis should be appropriately investigated to identify treatable causes and referral to a neurologist should be made so that appropriate neurological investigations can be performed (Grade C recommendation; level IV evidence)**
- **The severity of neuropathy should be sequentially graded with a validated tool, such as the Total Neuropathy Score. (Grade C recommendation; level IV evidence)**
- **Potentially neurotoxic drug treatments should be used with caution in patients with a pre-existing peripheral neuropathy (Grade C recommendation; level IV evidence)**
- **Any patient who develops a significant (e.g. >NCI grade 2) or progressive chemotherapy induced peripheral neuropathy should be managed with graded dose reduction or drug withdrawal. Referral to a neurologist for specialist investigation should be considered if there is no improvement or the use of another neurotoxic drug is planned (Grade C recommendation; level IV evidence).**
- **All patients with chronic peripheral neuropathic pain should be considered for multimodal analgesic treatment including an opioid, ion channel blocker and SNRI (Grade A recommendation; level 1b evidence).**
- **Superficial neuropathic pain should be treated with topical lidocaine 5% plaster (Grade A recommendation; level 1b evidence).**

- **Patients with significant uncontrolled neuropathic pain should be referred promptly for specialist advice regarding pain management (Grade C recommendation; level IV evidence)**
- **Routine use of nutritional supplements cannot be recommended due to insufficient evidence for benefit, although deficiencies should be corrected (Grade C recommendation; level IV evidence).**

7. Control of other symptoms

7.1 Nausea and vomiting

7.1.1 Acute emesis

Nausea and vomiting may occur in myeloma patients for many reasons – as symptoms of hypercalcaemia, side-effects of analgesics and toxicities of anti-cancer therapies. Acute onset emesis can usually be managed with dopaminergic prokinetic anti-emetics, eg oral domperidone or oral/parenteral metoclopramide. These have minimal side-effects, whereas the anticholinergic cyclizine may be effective but is associated with significant sedation and dry mouth. Note that whereas prokinetics stimulate GI motility, cyclizine suppresses it and so it is irrational to use a combination of these simultaneously.

Myeloma patients may be psychologically stressed by the illness and coping with treatment: stress-related hyperacidity and peptic ulceration may occur and be expressed by nausea and vomiting. This is best managed by an oral proton pump inhibitor (PPI).

If emesis does not settle with a dopamine agonist and PPI, then the butyrophenone drug haloperidol is usually effective. At doses of, for example, 1.5-5mg nocte as a single oral or sc dose (+ 0.5-2mg prn doses) it is usually not sedating. Older people may be more prone to extrapyramidal adverse effects.

If haloperidol is also unsuccessful, then the broad spectrum phenothiazine levomepromazine (Nozinan) may be used. When given at doses of up to 10mg/day (as a single nocte dose or by 24 hour infusion via a sc syringe driver), it is fairly well tolerated – but at higher doses it tends to be sedating, causes hypotension and may be associated with Parkinsonian side-effects. Levomepromazine should therefore be reserved for patients who do not respond to domperidone/metoclopramide or haloperidol.

Acute emesis induced by chemotherapy or radiation therapy can also be managed by agents such as domperidone or metoclopramide, but for the more intensive chemotherapy regimens used in myeloma serotonin (5HT₃) receptor antagonists may be considered (Keeley 2009). Local policies will determine which 5HT₃ receptor antagonist is used.

Prolonged use of these agents cause troublesome constipation, so they are best reserved for short-term use related to anti-cancer therapies.

Most of the regimens used in myeloma are not 'highly emetogenic' but some patients experience intolerable nausea and vomiting with even less emetogenic combinations. In such cases, the Neurokinin-1 (NK-1) antagonist aprepitant (Emend) may be used. Aprepitant has to be started, together with dexamethasone, on the day before chemotherapy. It is licensed for highly emetogenic chemotherapy or moderately emetogenic chemotherapy with anthracycline/cyclophosphamide combinations. It is reasonable to consider its use off licence for other patients who may otherwise refuse further chemotherapy (Kris *et al*, 2006).

7.1.2 Chronic emesis

Nausea and vomiting which does not settle even with the combinations described above may be managed with continuous subcutaneous infusions, delivered by a portable syringe driver, of a combination of metoclopramide and haloperidol, or metoclopramide and levomepromazine. A serotonin (5HT₃) receptor antagonist may be added for a few days but prolonged use will cause severe constipation. For example, the dose of levomepromazine may be increased up to 25mg/day, but significant sedation is likely result at this dose. At this level the patient should be referred to a palliative care team. It is important to exclude raised intracranial pressure and other causes for refractory emesis.

7.2 Loss of appetite (anorexia)

Classic cancer cachexia occurs frequently in patients with solid tumours but is uncommon in myeloma. However, many patients do experience periods of loss of appetite and may lose weight. An exceptional case is at the time of high dose chemotherapy and stem cell transplant, when significant weight loss may occur in a short time (Murray and Pindoria 2009). Often anorexia is related to stress and psychological adjustment disorders – referral to a specialist nurse for exploration of these issues may be the best first step. A good dietary history should be taken, preferably by a dietician. Patients should be encouraged to take smaller, more frequent meals rather than two large meals per day. Nutritional supplements can be added according to the patient's taste, eg milk-based

drinks, milkshakes or powders to increase the protein/carbohydrate content of savoury dishes.

Corticosteroids can improve appetite over a few days, but chronic use should be avoided because of their long-term adverse effects. For example, it is reasonable to give dexamethasone 4mg/day for a maximum of 7 days, tapering off over one week. For more prolonged appetite stimulation, the progestogen megestrol acetate can be used at doses of 160mg bd or tid. At these doses the side-effects of ankle oedema and DVT are minimal (Berenstein and Ortiz 2005).

7.3 Constipation and diarrhoea

Constipation occurs with great frequency in myeloma patients. It may be a result of hypercalcaemia, or a side-effect of drug therapy with thalidomide, opioids, anticholinergics or serotonin (5HT₃ receptor) antagonists. It is important to review all current medication and minimise drugs which may be causing or aggravating constipation.

Opioid-induced constipation can be reduced by using fentanyl or buprenorphine which cause much less of this adverse effect, compared to morphine and oxycodone.

Laxative medications are very commonly used but the evidence base for them is not convincing (Ahmedzai and Boland 2007). It is rational to consider the use of a combination of a stool softener such as docusate together with a stimulant such as senna or codanthrusate. For prevention of constipation, regular macrogol (Movicol, Laxido) or lactulose may be considered.

The management of opioid-induced constipation is likely to be revolutionised in the next few years with the advent of peripherally acting opioid antagonists (Holzer *et al*, 2009). These novel approaches include slow-release naloxone (available in a combination with slow release oxycodone –Targinact (Schutter *et al*, 2010); and parenteral methylnaltrexone (Thomas *et al*, 2008).

Patients receiving treatment for myeloma may suffer with diarrhoea. Bortezomib and lenalidomide may be associated with diarrhoea and require dose modification or cessation of the drug (Bird et al 2010). Management of diarrhoea should be directed at the underlying cause, but symptomatic measures include anti-motility agents such as loperamide, and anticholinergic drugs such as propantheline (Twycross and Wilcock 2008).

7.4 Mucositis

Oropharyngeal mucositis occurs frequently in myeloma patients on chemotherapy agents. It is most troublesome during the neutropenic phase of haemopoietic stem cell transplantation (HSCT), when it can lead to complete inability to take oral food or fluids because of the pain and ulceration.

Mucositis may be monitored and measured using a standardised visual rating tool. During HSCT this would reasonably be done at least once a day. Potential management for mild grades of mucositis includes the weak analgesic Difflam mouthwash. There is no place for chlorhexidine mouthwash. More troublesome cases can be managed with oxetacaine/antacid combination, which is swilled round the mouth and then may be swallowed to treat oropharyngeal and oesophageal mucositis. This may have to be given up to 2 hourly in grade 4 cases. Some patients can tolerate lidocaine ice lollies or mouthwashes (Peterson *et al*, 2009; Potting *et al*, 2006; Shaiova *et al*, 2004).

In patients who are at special risk of mucositis, the keratinocyte growth factor palifermin may be used, but it has to be started before chemotherapy (Kobbe *et al*, 2010). Caphasol may also be effective in the setting of high dose therapy (Papas et al 2003).

With grade 1-2 mucositis, the patient may be able to take oral medication such as an opioid tablet or liquid. Oral transmucosal fentanyl may be effective (Shaiova *et al*, 2004), but many patients with mucositis cannot use this. With more severe grades, a subcutaneous infusion of oxycodone (or morphine) can be given by syringe driver. In the most severe cases, it may be necessary to start an intravenous patients controlled analgesia (PCA) delivery of opioid – this should be done by the local acute pain team.

7.5 Sedation

Many drugs used in myeloma can cause sedation (thalidomide, opioids, gabapentin, tricyclics, benzodiazepines). In addition, drowsiness may be a feature of hypercalcaemia or of hypoactive delirium. It is important to warn patients of likely sedation when starting new medication and they should be counselled about the risks of taking sedative medication and drinking or driving .

Sedation due to the initiation of opioids sometimes settles in a few days, but often it does not. Fentanyl and oxycodone are less sedating than morphine or other potent opioids and the patient may become more alert within days of changing to these (Ahmedzai and Brooks 1997; Clark *et al*, 2004; Mucci-LoRusso *et al*, 1998; Reid *et al*, 2006).

If sedation is causing impairment of quality of life, then the patient should be referred to a palliative care team for consideration of a psychostimulant such as methylphenidate or modafanil (Blackhall *et al*, 2009; Gagnon *et al*, 2005; Minton *et al*, 2008).

7.6 Fatigue

Fatigue occurs in the majority of patients with myeloma and may be a major cause of reduced functioning and quality of life, although is often under-recognised by professionals. It is usually multifactorial – common treatable causes include anaemia, low testosterone, thyroid deficiency and there may be biochemical abnormalities including secretion of cytokines (Shafqat *et al*, 2005). Often the fatigue is related to sedating medication (see section above). Psychological causes of mental fatigue are common and should be explored by a specialist nurse or in severe cases, by a psychologist or psychiatrist.

There is clinical trial evidence that aerobic exercise programmes and the encouragement of physical activity after bouts of hospitalisation, especially for HSCT, are beneficial in terms of mood and physical functioning (Coleman *et al*, 2003; Dimeo *et al*, 1997). Patients should be encouraged to take brisk walks and graded exercises, including swimming. Anorexic patients may improve if given high carbohydrate supplements to boost calorie intake and the state of hydration should be reviewed. It is important to pay attention to sleep patterns, to encourage activity and minimal napping by day, and the avoidance of

getting up at night to take refreshments or watch television. Patients should be referred to physiotherapy or rehabilitation services for these programmes.

8. Bisphosphonate-induced Osteonecrosis of the Jaw (BONJ)

Osteonecrosis of the jaw (ONJ) is an increasingly recognized complication of BP therapy and is sometimes called Bisphosphonate induced osteonecrosis of the jaws (BONJ). The first case was described in 2003 (Carter and Goss 2003) The definition of BONJ is ‘ the presence of exposed necrotic bone in the mandible or maxilla that does not heal after 2 months in a patient receiving or with prior exposure to a bisphosphonate who has not had radiotherapy.’ Typical features are pain and localised infection, loosening of teeth and even spontaneous avulsion and soft tissue ulceration with sinus formation.

The precise aetiology is unclear but is likely to be multifactorial with contributions from a number of factors including infection and vascular insufficiency within the microcirculation. The most common precipitating event is dental extraction. A history of surgical intervention, long term use bisphosphonates especially sequential use of intravenous pamidronate/zoledronic therapy, prior history of malignant disease, significant comorbidity, smoking and poor dental health are all recognized risk factors (Boonyapakorn *et al*, 2008), there is also evidence of possible genetic risk factors (Sarasquete *et al*, 2009). Duration of therapy and dosage (exposure) seems to be an important aspect (Mavrokokki *et al*, 2007). The incidence of BONJ with oral bisphosphonates is much less than those administered intravenously (possibly 100 times less). With oral agents, the incidence of BONJ is estimated to be 1:10,000 and 1: 100,000 patient years (Khosla *et al*, 2007).

The diagnosis of BONJ is a clinical one. Patients present with pain and evidence of local infection (swelling, pus) although in a minority they are asymptomatic. The lesions are mostly localized to the mandible and over half occur in regions of recent dental extraction. Plain radiographs (OPG panoramic x-rays) can miss the presence of osteomyelitis in the early stages. A biopsy is seldom necessary as are further investigations such as CT, MRI or PET imaging.

The management of BONJ is largely supportive. Symptom control is usually achieved by good oral care with without antibiotics. Occasionally limited debridement of bone is indicated. There is no consensus as to whether to stop temporarily or discontinue bisphosphonate usage in affected individuals. Patients may develop BONJ up a year after

stopping BP treatment. Because BONJ tends to be a chronic condition, prevention is paramount and patients at risk should be identified and invasive dental procedures avoided (Barker and Rogers 2006; Weitzman *et al*, 2007).

Patients should be made aware of the risk of BONJ before embarking on long term BP use. Patients on oral bisphosphonates should be encouraged to see a dentist, however for those commencing intravenous bisphosphonates dental screening is essential (Weitzman *et al*, 2007). Before initiation of IV BP therapy the patient should have no expected extractions for the foreseeable future. Any teeth that are likely to require extraction should be removed before starting therapy. Ongoing dental care and maintenance (fillings) can continue with patients on the IV treatment. Patients, who have already received IV therapy and develop dental problems, which might possibly require extractions, should be referred to a specialist. It is important that teeth/roots are retained whenever this is feasible. Although there is little evidence to guide management of bisphosphonate treatment around invasive dental work it would seem reasonable to stop treatment in high risk patients and not recommence treatment until healing has occurred but this approach should be balanced against any ongoing fracture risks due to disease in the individual patient (Badros *et al*, 2008).

Recommendations

- **All patients to be started on long term bisphosphonate treatment should be warned of the risk of BONJ and its predisposing factors (Grade C recommendation; level IV evidence).**
- **All patients to be started on IV BP should be referred for a dental opinion and any teeth of poor prognosis extracted before initiation of BP therapy. Patients on long-term oral bisphosphonates should have regular dental care and maintain excellent oral hygiene (Grade C recommendation; level IV evidence).**
- **Invasive dental procedures in patients on IV or long-term oral bisphosphonate should be avoided as far as possible. For patients on IV BP, a specialist opinion should be sought prior to any extractions (Grade C recommendation; level IV evidence).**

- **Patients with suspected BONJ should be referred to a clinician with special interest and expertise in the management of this condition (Grade C recommendation; level IV evidence).**

9. The role of complementary therapies in the management of myeloma

9.1 Definition

Complementary therapy can be defined as therapies which are used alongside, or integrated with, conventional health care (Tavares 2003). These differ from alternative therapies, which are designed to be used in place of conventional therapy. However, a clear definition of what constitutes complementary and alternative medicine has not yet been elucidated, and therefore discretion must be exercised when interpreting guidance pertaining to these therapies.

In the UK, research is emerging on the role of complementary therapy in the management of myeloma. To date, the majority of available evidence is anecdotal and policy development is in its infancy. Studies have revealed greater than 25% of patients with haematological malignancies including myeloma, are using complementary therapy as adjunct to their prescribed conventional treatment (Molassiotis *et al*, 2005a). delivered primarily in hospices, hospitals and throughout the voluntary sector (Tavares 2003) .

Complementary therapy has a role in the management of multiple myeloma when used as adjunct to conventional medicine. It improves patients' perceived quality of life and ability to cope with the effects of the disease. The development of an evidence-base to support complementary therapy use in myeloma is in the early stages of development. Until complementary therapy has undergone rigorous testing in clinical trials, healthcare professionals should exercise caution with their recommendations.

9.2 The patient's preference

Patients with myeloma may express preference for complementary therapy and place value in the role they have to play within the context of their cancer care plan - for the management of both the psycho-social and physiological effects associated with myeloma. Patients may value complementary therapy and the sense of control gained when they are used as part of their cancer treatment plan. Consequently patient choice should be informed and respected by healthcare professionals in order to ensure the best overall treatment and care plan for myeloma is delivered.

Healthcare professionals should recognise that within the myeloma care plan, patient treatment preference, choice and need for reliable information is critical such that complementary therapy is accessed safely, should a patient wish to do so. In line with the NICE recommendations for Improving Supportive and Palliative Care for Adults with Cancer (2003b), the healthcare community must work within their Cancer Network to ensure availability of high quality information for complementary therapy, where there is evidence to support its use.

9.3 Potential benefits of complementary therapy

There is a dearth of scientific evidence to support the effectiveness of complementary therapy in the management of myeloma; however, some studies have shown that complementary therapy can help patients with myeloma to: manage their symptoms, live with altered body image, promote relaxation, alleviate anxiety, reduce chemotherapy side-effects, improve sleep pattern, reduce stress and tension, reduce psychological distress / provide emotional support and improve well-being (Molassiotis *et al*, 2005a), Leukemia and Lymphoma Society (2006). Importantly, cancer patients using complementary therapy also perceive an improved quality of life (Tovey *et al*, 2005).

Some complementary therapies, such as acupuncture, have been submitted to more rigorous evaluation and are acknowledged for their effective use in cancer treatment for the management of chemotherapy-associated nausea and vomiting (1997). However, no convincing scientific-evidence has emerged to date that shows complementary therapy slows cancer progression (2006).

9.4 Commonly used complementary therapies

The types of complementary therapies and frequency with which they are used by myeloma patients vary considerably. Among the most common therapies are homeopathy, touch therapies such as aromatherapy, massage and reflexology, healing and energy therapies such as reiki, spiritual healing and therapeutic touch, hypnosis and hypnotherapy, acupuncture, herbal medicines and dietary interventions (Molassiotis *et al*, 2005b) . Examples of the latter include green tea and curcumin, two chemoprotective agents shown to induce cellular apoptosis in myeloma (Bharti *et al*, 2003; Hatcher *et al*, 2008; Liu *et al*, 2007; Nakazato *et al*, 2005; Shamma *et al*, 2006) and currently gaining credibility as effective complementary therapies for myeloma. In addition, as covered in section 6.4, some topical treatments have a physiological basis in providing symptomatic relief in painful peripheral neuropathy (Vriens *et al* 2008). However, caution should be exercised with some nutritional supplements. For example, there is evidence that nutritional supplements containing ascorbic acid (vitamin C) may inhibit the anti-myeloma effect of bortezomib and should be avoided at least on the days of bortezomib administration (Peronne *et al* 2009).

Recommendations

- **Complementary therapy can offer benefit to myeloma patients when integrated with conventional medical treatments (Grade B recommendation; level IIIb evidence)**
- **Patients, carers and healthcare professionals should have access to high-quality information on the role of complementary therapy in the management of myeloma. Reputable support organisations, such as Myeloma UK, are a potential source of this information (Grade C recommendation; level IV evidence)**
- **Healthcare professionals should be aware of and prepared to appraise their patient's choice and preference for complementary therapy (Grade C recommendation; level IV evidence)**
- **Careful consideration should be applied by healthcare professionals before recommending complementary therapy until a rigorous evidence-base is developed (Grade C recommendation; level IV evidence).**

10. End of life care

10.1 Definitions

End of life care has recently been the subject of a major review by the NHS, leading to an end of life strategy by the Department of Health Report (2008). It is likely that all SHAs, PCTs and acute trusts will be undertaking regional and local reviews of their end of life care provision. Local policies and guidelines are being drawn up and so this section will cover the main concepts only.

Definitions relating to the 'end of life' vary. The 'end of life' strategy and the Darzi report apply to patients in their last months or even years of life, but they do not provide unequivocal definitions. The 'end of life care pathway' (eg Liverpool EOL CP) comes into action when the individual is thought to have 24-48 hours prognosis. It is easy for these terms to be misapplied and there is a possibility for patients and carers to be upset by referral to the wrong agency.

10.2 Recognition of end of life

In myeloma care, it is important for clinicians and teams to recognise when a patient has advancing and untreatable disease to the point that death is likely to occur within the next twelve months. Usually this is evident by relapse which is confirmed by investigations including bone marrow, blood counts and protein levels, and imaging including MRI of skeletal axis, At this point, which should be a joint decision of the MDT, it is reasonable to refer the patient to a palliative care team if this has not already been done. This will allow for a palliative care team member to become acquainted with the patient even if he or she does not have significant symptom needs at the present. A holistic needs assessment by the referring team or by the palliative care team, can highlight new problems and issues apart from symptoms, e.g. psychosocial difficulties, existential concerns and decisions regarding future care.

One of the UK government's end of life programmes is about 'Preferred Priorities for Care' (PPC). This includes eliciting the patients and carers' views and preferences for where the remaining care should be (i.e. home, hospital or hospice, or often a combination); what treatments the patient would want to have or avoid having; and the patient's views on cardiopulmonary resuscitation.

The results of these discussions should be conveyed to the GP, as patients with advanced cancer are now being placed by GPs on their registers for further community-based follow-up, in accordance with the so-called 'Gold Standards Framework'.

10.3 Continuation of anti-cancer and other active therapies

The withdrawal of anti-cancer treatment may be seen by many patients as a severe blow to morale and it is essential for this discussion to be held in the presence of family members and a specialist nurse. It should be followed up by a discussion of the patient's and carers' preferences for future care, including the place of care towards the end of life and the use of life-maintaining therapies.

In myeloma management, it is possible to keep offering the patient further forms of treatment aimed at reducing bone marrow plasma cell activity and consequent expression of myeloma symptoms. Even when the patient is approaching the terminal stage (last weeks of life) and specific anti-cancer treatments have been withdrawn, blood and platelet transfusions can be helpful in maintaining quality of life by relieving exertional dyspnoea and preventing bleeding. Management of hypercalcaemia and the maintenance of hydration can be very helpful to prevent symptoms and the onset of renal failure. In occasional patients, management of symptomatic hyperviscosity with regular plasma exchange may be reasonable, and may permit safe transfusion of blood.

If the patient is referred to a palliative care service at this stage, it is helpful for the haematology team to continue to share the care by offering admissions for blood or platelets and other more invasive measures, which may be problematic in a palliative care setting such as many hospices. However, after each blood or platelet transfusion at this stage, it is important for the patient to be asked how worthwhile it has been, as a point may be reached when the patient would actually prefer not to have further transfusions as the benefits do not outweigh the difficulties associated with hospital admission.

At this stage, the balance of responsibility for care may change significantly between GP, hospital and community palliative care teams, although the haematology team, and the patient's clinical nurse specialist (CNS) in particular, often provide continuity of care.

10.4 Last days of life

The term 'last days of life' is preferred to 'end of life' at this stage, as it focuses the attention of the team and the family on the patient's main priorities (and also, as mentioned above, 'end of life' can also refer to the last years of life in chronic disease). This stage is reached when the patient has become increasingly bedbound, profoundly anorexic, weak and spending increasing time at reduced levels of consciousness. Pain and other symptoms may become refractory to previously satisfactory management, in which case a palliative care team should be called upon if the patient is in hospital. It should be noted that family carers and dependents have their own needs at this stage of illness and the haematology CNS, together with the palliative care team, social services and primary care team, should look for and address these needs. Therefore both the patient's and family's holistic needs have to be screened at regular intervals once the patient enters the terminal stage (2004b).

Specific issues which need attention in the last days of life include –

- Initiation of the 'end of life' (better called 'last days of life') pathway. Most hospitals should now have a policy on how to implement this process, which is a quality assurance tool to ensure that inappropriate medications and obtrusive observations are stopped and all important symptoms are covered by regular or prn prescribing. The pathway also checks if the patient's wishes for a final place of care and other preferences are being met, where possible. Very importantly, the pathway also helps the caring team to recognise the needs of family members for information, emotional and social support and can prepare the way for bereavement support for those who need it.
- For prn medication, a subcutaneous butterfly cannula may be placed to avoid skin punctures
- For regular medication, one or more syringe drivers may be started to give drugs that can be delivered subcutaneously.
- Fentanyl or buprenorphine patches, if already started, should preferably be left in place, and additional pain medication given by oral or subcutaneous route.

- For patients who become unconscious and start to make the 'death rattle', this can be alleviated by positioning and judicious suctioning to remove pharyngeal (not tracheal) mucus, while injections or an infusion of an anticholinergic drug, such as hyoscine butylbromide (Buscopan) or glycopyrrolate should be started to prevent further mucus secretion.
- The roles and needs of family and others close to the patient require recognition and support.

11. Conclusions – the 'holistic needs assessment' and clinical research parameters

In conclusion, patients with myeloma not only require treatments directed at their disease activity but also a wide range of supportive and palliative measures to optimise quality of life at all stages of their disease. Most aspects of myeloma management can be addressed and delivered by the treating haematologists, but collaboration may be required from other relevant specialists. The demands of effective supportive and palliative management are variable between patients and present significant challenges for treating haematologists and other health professionals. Levels of unmet supportive care needs have been identified in a quarter of patients with myeloma and a third of their partners, with significant components of anxiety and depression in both groups (Molassiotis *et al*, 2010).

The UK NHS NICE Guidance for Supportive and Palliative Care for adults with cancer (2004b) recommends that all cancer patients should have an 'holistic needs assessment' at key points in their trajectory, i.e. at diagnosis, after initial treatment, during follow-up. The specification for how to measure holistic needs in cancer patients was produced by the Kings College Department of Cancer Nursing (Richardson 2006). One useful instrument for this purpose is the 'SPARC' tool, which is completed by the patient. It identifies and scores needs with respect to uncontrolled pain and other symptoms as well as psychosocial, spiritual, financial and other issues, using a 4-point verbal or Yes/No rating scale (Ahmed *et al*, 2009). An alternative approach is the Distress Thermometer, which rates overall distress on a numerical analogue scale (Ransom *et al*, 2006).

The increasing survival of myeloma patients and associated symptom burden raises the need for clinical research to run in parallel to novel approaches to controlling disease. Patients in clinical trials will usually need a formal tool for assessing symptoms and their

response to treatments. The recommended method is use of the EORTC QLQ-C30 generic quality of life questionnaire, together with its myeloma-specific module (QLQ-MY20 (Cocks *et al*, 2007). A new instrument has been developed by the EORTC for the measurement of pain arising from bone metastases (Chow *et al*, 2009) which may be useful in some settings. Lessons learnt from the field of late effects of cancer and its treatments (Eiser *et al*, 2007), conventionally applied to curable malignancy, may be of increasing importance in the ongoing management of long term survivors of myeloma.

Recommendations

- **All patients should be assessed for unmet holistic needs at key times in the disease trajectory, preferably using a formal patient-rated tool such as SPARC or the Distress Thermometer (Grade C recommendation; level IV evidence)**
- **For research studies, the EORTC family of quality of life instruments should be used (Grade B recommendation; level IIb evidence)**
- **The recommendations in these guidelines based on level II-IV evidence identify areas where there is scope for future research (Grade C recommendation; level IV evidence)**

12: Levels of evidence and grades of recommendation

Levels of Evidence

Ia	Evidence obtained from meta-analysis of randomised controlled trials
Ib	Evidence obtained from at least one randomised controlled trial
IIa	Evidence obtained from at least one well-designed, non-randomised study, including phase II trials and case-control studies
IIb	Evidence obtained from at least one other type of well-designed, quasi-experimental study, i.e. studies without planned intervention, including observational studies
III	Evidence obtained from well-designed, non-experimental descriptive studies. Evidence obtained from meta-analysis or randomised controlled trials or phase II studies which is published only in abstract form
IV	Evidence obtained from expert committee reports or opinions and/or clinical experience of respected authorities

Grades of Recommendation

Grade A Evidence level Ia, Ib	Recommendation based on at least one randomised controlled trial of good quality and consistency addressing specific recommendation
Grade B Evidence level IIa, IIb, III	Recommendation based on well conducted studies but no randomised controlled trials on the topic of recommendation
Grade C Evidence level IV	Evidence from expert committee reports and/or clinical experiences of respected authorities

13. Declarations of Interest

All the authors have filled in a declaration of interest from which has been reviewed by the chair of the BCSH Haemato Oncology Task Force and the chair of BCSH and none of the authors are considered to have a conflict of interest with regards to the recommendations made in this guideline.

REFERENCES

- (1997) Acupuncture. NIH Consensus Statement. <http://consensus.nih.gov/1997/1997acupuncture107html.htm>. **15**, 1-34.
- (2003a) National Institute for Clinical Excellence. Balloon kyphoplasty for vertebral compression fractures. *Interventional Procedure Guidance*, **20**.
- (2003b) National Institute for Health and Clinical Excellence. Improving outcomes in haematological cancers-The Manual. <http://www.nice.org.uk/docref.asp?d=90429>.
- (2004a) Guidelines on the diagnosis and management of AL amyloidosis. Guidelines Working Group of UK Myeloma Forum; British Committee for Standards in Haematology, British Society for Haematology. *British Journal of Haematology*, **125**, 681-700.
- (2004b) National Institute for Clinical Excellence Guidance on Cancer Services. Improving Supportive and Palliative Care for Adults with Cancer - The Manual <http://www.nice.org.uk/nicemedia/live/10893/28816/28816.pdf>.
- (2006) Leukemia and Lymphoma Society. Integrative Medicine and Complementary and Alternative Therapies as part of Blood Cancer Care. http://www.leukemia-lymphoma.org/attachments/National/br_1150734030.pdf.
- (2008) Department of Health End of Life Care Strategy. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_086277.
- Ahmed, N., Bestall, J.C., Payne, S.A., Noble, B. & Ahmedzai, S.H. (2009) The use of cognitive interviewing methodology in the design and testing of a screening tool for supportive and palliative care needs. *Supportive Care in Cancer*, **17**, 665-673.
- Ahmedzai, S. & Brooks, D. (1997) Transdermal fentanyl versus sustained-release oral morphine in cancer pain: preference, efficacy, and quality of life. The TTS-Fentanyl Comparative Trial Group. *Journal of Pain and Symptom Management*, **13**, 254-261.
- Ahmedzai, S.H. & Boland, J. (2007) The total challenge of cancer pain in supportive and palliative care. *Current Opinion in Supportive and Palliative Care*, **1**, 3-5.
- Argyriou, A.A., Iconomou, G. & Kalofonos, H.P. (2008) Bortezomib-induced peripheral neuropathy in multiple myeloma: a comprehensive review of the literature. *Blood*, **112**, 1593-1599.
- Badros, A., Goloubeva, O., Dalal, J.S., Can, I., Thompson, J., Rapoport, A.P., Heyman, M., Akpek, G. & Fenton, R.G. (2007) Neurotoxicity of bortezomib therapy in multiple myeloma: a single-center experience and review of the literature. *Cancer*, **110**, 1042-1049.
- Badros, A., Terpos, E., Katodritou, E., Goloubeva, O., Kastiris, E., Verrou, E., Zervas, K., Baer, M.R., Meiller, T. & Dimopoulos, M.A. (2008) Natural history of osteonecrosis of the jaw in patients with multiple myeloma. *Journal of Clinical Oncology*, **26**, 5904-5909.
- Ballantyne, J.C. & Mao, J. (2003) Opioid therapy for chronic pain. *New England Journal of Medicine*, **349**, 1943-1953.
- Barker, K. & Rogers, S. (2006) Bisphosphonate-associated Osteonecrosis of the jaws: a guide for the general dental practitioner. *Dental Update*, **33**, 270-275.
- Baron, F., Frere, P. & Beguin, Y. (2003) Once weekly recombinant human erythropoietin therapy is very efficient after allogeneic peripheral blood stem cell transplantation when started soon after engraftment. *Haematologica*, **88**, 718-720.
- Baron R, Mayoral V, Leijon G, Binder A, Steigerwald I, Serpell M (2009). 5% lidocaine medicated plaster versus pregabalin in post-herpetic neuralgia and diabetic

- polyneuropathy: an open-label, non-inferiority two-stage RCT study. *Curr Med Res Opin.*; **25**:1663-76
- Baz, R., Li, L., Kottke-Marchant, K., Srkalovic, G., McGowan, B., Yiannaki, E., Karam, M.A., Faiman, B., Jawde, R.A., Andresen, S., Zeldis, J. & Hussein, M.A. (2005) The role of aspirin in the prevention of thrombotic complications of thalidomide and anthracycline-based chemotherapy for multiple myeloma. *Mayo Clinic Proceedings*, **80**, 1568-1574.
- Bennett, C.L., Angelotta, C., Yarnold, P.R., Evens, A.M., Zonder, J.A., Raisch, D.W. & Richardson, P. (2006) Thalidomide- and lenalidomide-associated thromboembolism among patients with cancer. *Journal of the American Medical Association*, **296**, 2558-2560.
- Bennett, M. (2001) The LANSS Pain Scale: the Leeds assessment of neuropathic symptoms and signs. *Pain*, **92**, 147-157.
- Bennett, M.I., Attal, N., Backonja, M.M., Baron, R., Bouhassira, D., Freynhagen, R., Scholz, J., Tölle, T.R., Wittchen, H.U. & Jensen, T.S. (2007) Using screening tools to identify neuropathic pain. *Pain*, **127**, 199-203.
- Berenstein, E.G. & Ortiz, Z. (2005) Megestrol acetate for the treatment of anorexia-cachexia syndrome. *Cochrane Database Systemic Reviews*, **18**, CD004310.
- Bharti, A.C., Donato, N., Singh, S. & Aggarwal, B.B. (2003) Curcumin (diferuloylmethane) down-regulates the constitutive activation of nuclear factor-kappa B and I κ B kinase in human multiple myeloma cells, leading to suppression of proliferation and induction of apoptosis. *Blood*, **101**, 1053-1062.
- Binder A, Bruxelles J, Rogers P, Hans G, Bösl I, Baron R (2009). Topical 5% lidocaine (lignocaine) medicated plaster treatment for post-herpetic neuralgia: results of a double-blind, placebo-controlled, multinational efficacy and safety trial. *Clin Drug Investig.* **29**:393-408
- Bird, J., Behrens, J., Westin, J., Turesson, I., Drayson, M., Beetham, R., D'Sa, S., Soutar, R., Waage, A., Gulbrandsen, N., Gregersen, H. & Low, E. (2009) UK Myeloma Forum (UKMF) and Nordic Myeloma Study Group (NMSG): guidelines for the investigation of newly detected M-proteins and the management of monoclonal gammopathy of undetermined significance (MGUS). *British Journal of Haematology*, **147**, 22-42.
- Birgegard, G., Gascon, P. & Ludwig, H. (2006) Evaluation of anaemia in patients with multiple myeloma and lymphoma: findings of the European CANCER ANAEMIA SURVEY. *European Journal of Haematology*, **77**, 378-386.
- Blackhall, L., Petroni, G., Shu, J., Baum, L. & Farace, E. (2009) A pilot study evaluating the safety and efficacy of modafinil for cancer-related fatigue. *Journal of Palliative Medicine*, **12**, 433-439.
- Boggio, L. & Green, D. (2001) Recombinant human factor VIIa in the management of amyloid-associated factor X deficiency. *British Journal of Haematology*, **112**, 1074-1075.
- Bohlius, J., Schmidlin, K., Brillant, C., Schwarzer, G., Trelle, S., Seidenfeld, J., Zwahlen, M., Clarke, M., Weingart, O., Kluge, S., Piper, M., Rades, D., Steensma, D.P., Djulbegovic, B., Fey, M.F., Ray-Coquard, I., Machtay, M., Moebus, V., Thomas, G., Untch, M., Schumacher, M., Egger, M. & Engert, A. (2009) Recombinant human erythropoiesis-stimulating agents and mortality in patients with cancer: a meta-analysis of randomised trials. *Lancet*, **373**.
- Bokemeyer, C., Aapro, M.S., Courdi, A., Foubert, J., Link, H., Osterborg, A., Repetto, L. & Soubeyran, P. (2007) EORTC guidelines for the use of erythropoietic proteins in anaemic patients with cancer: 2006 update. *European Journal of Cancer*, **43**, 258-270.

Boonyapakorn, T., Schirmer, I., Reichart, P.A., Sturm, I. & Massenkeil, G. (2008) Bisphosphonate-induced osteonecrosis of the jaws: prospective study of 80 patients with multiple myeloma and other malignancies. *Oral Oncology*, **44**, 857-869.

Brenner, H., Gondas, A. & Pulte, D. (2008) Recent major improvement in long-term survival of younger patients with multiple myeloma. *Blood*, **111**, 2521-2526.

Brenner, H., Gondas, A. & Pulte, D. (2009) Expected long-term survival of patients diagnosed with multiple myeloma in 2006-2010. *Haematologica*, **94**, 270-275.

Bringhen S, Larocca A, Rossi D, Cavalli M, Genuardi M, Ria R, Gentili S, Patriarca F, Nozzoli C, Levi A, Guglielmelli T, Benevolo G, Callea V, Rizzo V, Cangialosi C, Musto P, De Rosa L, Liberati AM, Grasso M, Falcone AP, Evangelista A, Cavo M, Gaidano G, Boccadoro M, Palumbo A. (2010) Efficacy and safety of once weekly bortezomib in multiple myeloma patients. *Blood*. Aug 31. [Epub ahead of print]

Caraceni, A., Zecca, E., Bonezzi, C., Arcuri, E., Yaya, T.R., Maltoni, M., Visentin, M., Gorni, G., Martini, C., Tirelli, W., Barbieri, M. & De Conno, F. (2004) Gabapentin for neuropathic cancer pain: a randomized controlled trial from the Gabapentin Cancer Pain Study Group. *Journal of Clinical Oncology*, **22**, 2909-2917.

Carter, G.D. & Goss, A.N. (2003) Bisphosphonates and avascular necrosis of the jaws. *Australian Dental Journal*, **48**, 268.

Cavaletti, G., Beronio, A., Reni, L., Ghiglione, E., Schenone, A., Briani, C., Zara, G., Cocito, D., Isoardo, G., Ciaramitaro, P., Plasmati, R., Pastorelli, F., Frigo, M., Piatti, M. & Carpo, M. (2004) Thalidomide sensory neurotoxicity: a clinical and neurophysiologic study. *Neurology*, **62**, 2291-2293.

Cavaletti, G., Frigeni, B., Lanzani, F., Piatti, M., Rota, S., Briani, C., Zara, G., Plasmati, R., Pastorelli, F., Caraceni, A., Pace, A., Manicone, M., Lissoni, A., Colombo, N., Bianchi, G. & Zanna, C. (2007) The Total Neuropathy Score as an assessment tool for grading the course of chemotherapy-induced peripheral neurotoxicity: comparison with the National Cancer Institute-Common Toxicity Scale. *Journal of the peripheral nervous system*, **12**, 210-215.

Cavaletti, G. & Nobile-Orazio, E. (2007) Bortezomib-induced peripheral neurotoxicity: still far from a painless gain. *Haematologia*, **92**, 1308-1310.

Cella, D., Kallich, J., McDermott, A. & Xu, X. (2004) The longitudinal relationship of hemoglobin, fatigue and quality of life in anemic cancer patients: results from five randomized clinical trials. *Annals of Oncology*, **15**, 979-986.

Chapel H.M., Lee M., Hargreaves R., Pamphilon D.H., Prentice A.G. (1994) Randomised trial of intravenous immunoglobulin as prophylaxis against infection in plateau-phase multiple myeloma. The UK Group for Immunoglobulin Replacement Therapy in Multiple Myeloma. *Lancet*, **343**, 1059-63.

Chow, E., Hird, A., Velikova, G., Johnson, C., Dewolf, L., Bezjak, A., Wu, J., Shafiq, J., Sezer, O., Kardamakis, D., Linden, Y., Ma, B., Castro, M., Arnalot, P., Ahmedzai, S., Clemons, M., Hoskin, P., Yee, A., Brundage, M. & Bottomley, A. (2009) The European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire for patients with bone metastases: the EORTC QLQ-BM22. *European Journal of Cancer*, **45**, 1146-1152.

Clark, A.J., Ahmedzai, S.H., Allan, L.G., Camacho, F., Horbay, G.L., Richarz, U. & Simpson, K. (2004) Efficacy and safety of transdermal fentanyl and sustained-release oral morphine in patients with cancer and chronic non-cancer pain. *Current Medical Research and Opinion*, **20**, 1419-1428.

Cocks, K., Cohen, D., Wisloff, F., Sezer, O., Lee, S., Hippe, E., Gimsing, P., Turesson, I., Hajek, R., Smith, A., Graham, L., Phillips, A., Stead, M., Velikova, G. & Brown, J. (2007) An international field study of the reliability and validity of a disease-specific

questionnaire module (the QLQ-MY20) in assessing the quality of life of patients with multiple myeloma. *European Journal of Cancer*, **43**, 1670-1678.

Coleman, E.A., Coon, S., Hall-Barrow, J., Richards, K., Gaylor, D. & Stewart, B. (2003) Feasibility of exercise during treatment for multiple myeloma. *Cancer Nursing*, **26**, 401-409.

Coleman, M., Vigliano, E.M., Weksler, M.E. & Nachman, R.L. (1972) Inhibition of fibrin monomer polymerization by lambda myeloma globulins. *Blood*, **39**, 210-223.

Davies, A.N., Dickman, A., Reid, C., Stevens, A.M. & Zeppetella, G. (2009) Science Committee of the Association for Palliative Medicine of Great Britain and Ireland. The management of cancer-related breakthrough pain: recommendations of a task group of the Science Committee of the Association for Palliative Medicine of Great Britain and Ireland. *European Journal of Pain*, **13**, 331-338.

Di Minno, G., Coraggio, F., Cerbone, A., Capitanio, A., Manzo, C., Spina, M., Scarpato, P., Dattoli, G.M.R., Mattioli, P.L. & Mancini, M. (1986) Myeloma Paraprotein with Specificity for Platelet Glycoprotein IIIa in a patient with a Fatal Bleeding Disorder. *Journal of Clinical Investigation*, **77**, 157-164.

Dimeo, F.C., Tilmann, M.H., Bertz, H., Kanz, L., Mertelsmann, R. & Keul, J. (1997) Aerobic exercise in the rehabilitation of cancer patients after high dose chemotherapy and autologous peripheral stem cell transplantation. *Cancer*, **79**, 1717-1722.

Dimopoulos, M., Spencer, A., Attal, M., Prince, H.M., Harousseau, J.L., Dmoszynska, A., San Miguel, J., Hellmann, A., Facon, T., Foà, R., Corso, A., Masliak, Z., Olesnyckyj, M., Yu, Z., Patin, J., Zeldis, J.B. & Knight, R.D. (2007) Lenalidomide plus dexamethasone for relapsed or refractory multiple myeloma. *New England Journal of Medicine*, **357**, 2123-2132.

Dispenzieri, A. & Gertz, M.A. (2004) Treatment of POEMS syndrome. *Current Treatment Options in Oncology*, **5**, 249-257.

Eiser, C., Absolom, K., Greenfield, D., Snowden, J., Coleman, R., Hancock, B. & Davies, H. (2007) Follow-up care for young adult survivors of cancer: lessons from pediatrics. *Journal of Cancer Survivorship*, **1**, 75-86.

El-Cheikh, J., Stoppa, A.M., Bouabdallah, R., de Lavallade, H., Coso, D., de Collela, J.M., Auran-Schleinitz, T., Gastaut, J.A., Blaise, D. & Mohty, M. (2008) Features and risk factors of peripheral neuropathy during treatment with bortezomib for advanced multiple myeloma. *Clinical Lymphoma & Myeloma*, **8**, 146-152.

Federici, A.B. (2006) Acquired von Willebrand syndrome: an underdiagnosed and misdiagnosed bleeding complication in patients with lymphoproliferative and myeloproliferative disorders. *Seminars in Hematology*, **43**, S48-58.

Fourney, D.R., Schomer, D.F., Nader, R., Chlan-Fourney, J., Suki, D., Ahrar, K., Rhines, L.D. & Gokaslan, Z.L. (2003) Percutaneous vertebroplasty and kyphoplasty for painful vertebral body fractures in cancer patients. *Journal of Neurosurgery*, **98**, 21-30.

Gagnon, B., Low, G. & Schreier, G. (2005) Methylphenidate hydrochloride improves cognitive function in patients with advanced cancer and hypoactive delirium: a prospective clinical study. *Journal of Psychiatry and Neuroscience*, **30**, 100-107.

Geerts, W.H., Pineo, G.F., Heit, J.A., Bergqvist, D., Lassen, M.R., Colwell, C.W. & Ray, J.G. (2004) Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*, **123**, 338S-400S.

Hatcher, H., Planalp, R., Cho, J., Torti, F.M. & Torti, S.V. (2008) Curcumin: from ancient medicine to current clinical trials. *Cellular and Molecular Life Sciences*, **65**, 1631-1652.

Hedenus, M., Adriansson, M., San Miguel, J., Kramer, M.H., Schipperus, M.R., Juvonen, E., Taylor, K., Belch, A., Altes, A., Martinelli, G., Watson, D., Matcham, J., Rossi, G. & Littlewood, T.J. (2003) Efficacy and safety of darbepoetin alfa in anaemic

patients with lymphoproliferative malignancies: a randomized, double-blind, placebo-controlled study. *British Journal of Haematology*, **122**, 394-403.

Ho, T.W., Backonja, M., Ma, J., Leibensperger, H., Froman, S. & Polydefkis, M. (2009) Efficient assessment of neuropathic pain drugs in patients with small fiber sensory neuropathies. *Pain*, **141**, 19-24.

Holdcroft, A. & Power, I. (2003) Recent developments: management of pain. *British Medical Journal*, **326**, 635-639.

Holzer, P., Ahmedzai, S.H., Niederle, N., Leyendecker, P., Hopp, M., Bosse, B., Spohr, I. & Reimer, K. (2009) Opioid-induced bowel dysfunction in cancer-related pain: causes, consequences, and a novel approach for its management. *Journal of Opioid Management*, **5**, 145-151.

Jagannath, S., Barlogie, B., Berenson, J.R., Singhal, S., Alexanian, R., Srkalovic, G., Orlowski, R.Z., Richardson, P.G., Anderson, J., Nix, D., Esseltine, D.L. & Anderson, K.C. (2005a) Bortezomib in recurrent and/or refractory multiple myeloma. Initial clinical experience in patients with impaired renal function. *Cancer*, **103**, 1195-1200.

Jagannath, S., Durie, B.G., Wolf, J., Camacho, E., Irwin, D., Lutzky, J., McKinley, M., Gabayan, E., Mazumder, A., Schenkein, D. & Crowley, J. (2005b) Bortezomib therapy alone and in combination with dexamethasone for previously untreated symptomatic multiple myeloma. *British Journal of Haematology*, **129**, 776-783.

Jensen, M.E. & Kallmes, D.E. (2002) Percutaneous vertebroplasty in the treatment of malignant spine disease. *Cancer Journal*, **8**, 194-206.

Jensen, T.S., Madsen, C.S. & Finnerup, N.B. (2009) Pharmacology and treatment of neuropathic pains. *Current Opinion in Neurology*, **22**, 467-474.

Kariyawasan, C.C., Hughes, D.A., Jayatillake, M.M. & Mehta, A.B. (2007) Multiple myeloma: causes and consequences of delay in diagnosis. *Quarterly Journal of Medicine*, **100**, 635-664.

Kearon, C., Kahn, S.R., Agnelli, G., Goldhaber, S., Raskob, G.E. & Comerota, A.J. (2008) American College of Chest Physicians. Antithrombotic therapy for venous thromboembolic disease: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest*, **133**, 454S-545S.

Keeley, P.W. (2009) Nausea and vomiting in people with cancer and other chronic diseases. *Clinical Evidence (Online)*, pii, 2406.

Kelly, J.J., Kyle, R.A., O'Brien, P.C. & Dyck, P.J. (1981) Prevalence of monoclonal protein in peripheral neuropathy. *Neurology*, **31**, 1480-1483.

Keskinbora, K., Pekel, A.F. & Aydinli, I. (2007) Gabapentin and an opioid combination versus opioid alone for the management of neuropathic cancer pain: a randomized open trial. *Journal of Pain and Symptom Management*, **34**, 183-189.

Khosla, S., Burr, D., Cauley, J., Dempster, D.W., Ebeling, P.R., Felsenberg, D., Gagel, R.F., Gilsanz, V., Guise, T., Koka, S., McCauley, L.K., McGowan, J., McKee, M.D., Mohla, S., Pendrys, D.G., Raisz, L.G., Ruggiero, S.L., Shafer, D.M., Shum, L., Silverman, S.L., Van Poznak, C.H., Watts, N., Woo, S.B. & Shane, E. (2007) Bisphosphonate-associated osteonecrosis of the jaw: report of a task force of the American Society for Bone and Mineral Research. *Journal of Bone and Mineral Research*, **22**, 1479-1491.

Kobbe, G., Bruns, I., Schroeder, T., Czibere, A., Warnecke, J., Hieronimus, N., Safaian, N., Kondakci, M., Saure, C., Germing, U., Haas, R. & Fenk, R. (2010) A 3-day short course of palifermin before HDT reduces toxicity and need for supportive care after autologous blood stem-cell transplantation in patients with multiple myeloma. *Annals of Oncology*, March 16, epub ahead of print.

Kris, M.G., Hesketh, P.J., Somerfield, M.R., Feyer, P., Clark-Snow, R., Koeller, J.M., Morrow, G.R., Chinnery, L.W., Chesney, M.J., Gralla, R.J. & Grunberg, S.M. (2006)

- American Society of Clinical Oncology guideline for antiemetics in oncology: update 2006. *Journal of Clinical Oncology*, **24**, 2932-2947.
- Kristinsson, S.Y., Fears, T.R., Gridley, G., Turesson, I., Mellqvist, U.H., Björkholm, M. & Landgren, O. (2008) Deep vein thrombosis after monoclonal gammopathy of undetermined significance and multiple myeloma. *Blood*, **112**, 3582-3586.
- Kyle, R.A. (1992) Monoclonal proteins in neuropathy. *Neurologic Clinics*, **10**, 713-734.
- Kyle, R.A. & Gertz, M.A. (1995) Primary systemic amyloidosis: clinical and laboratory features in 474 cases. *Semin Hematol*, **32**, 45-59.
- Kyle, R.A., Gertz, M.A., Witzig, T.E., Lust, J.A., Lacy, M.Q., Dispenzieri, A., Fonseca, R., Rajkumar, S.V., Offord, J.R., Larson, D.R., Plevak, M.E., Therneau, T.M. & Greipp, P.R. (2003) Review of 1027 patients with newly diagnosed multiple myeloma. *Mayo Clinic Proceedings*, **78**, 21-33.
- Lane, J.M., Hong, R., Koob, J., Kiechle, T., Niesvizky, R., Pearse, R., Siegel, D. & Poynton, A.R. (2004) Kyphoplasty enhances function and structural alignment in multiple myeloma. *Clinical Orthopaedic Related Research*, **426**, 49-53.
- Lee, A.Y., Levine, M.N., Baker, R.I., Bowden, C., Kakkar, A.K., Prins, M., Rickles, F.R., Julian, J.A., Haley, S., Kovacs, M.J. & Gent, M. (2003) Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. *New England Journal of Medicine*, **349**, 146-153.
- Leigh, B.R., Kurtt, T.A., Mack, C.F., Matzner, M.B. & Shimm, D.S. (1993) Radiation therapy for the palliation of multiple myeloma. *International Journal of Radiation Oncology, Biology, Physics*, **25**, 801-804.
- Lennernäs, B., Frank-Lissbrant, I., Lennernäs, H., Kälkner, K.M., Derrick, R. & Howell, J. (2010) Sublingual administration of fentanyl to cancer patients is an effective treatment for breakthrough pain: results from a randomized phase II study. *Palliative Medicine*, **24**, 286-293.
- Levine, S. & Saltzman, A. (2004) Pyridoxine (vitamin B6) neurotoxicity: enhancement by protein-deficient diet. *Journal of Applied Toxicology*, **24**, 497-500.
- Littlewood, T.J., Bajetta, E., Nortier, J.W., Vercaemmen, E. & Rapoport, B. (2001) Effects of epoetin alfa on hematologic parameters and quality of life in cancer patients receiving nonplatinum chemotherapy: results of a randomized, double-blind, placebo-controlled trial. *Journal of Clinical Oncology*, **19**, 2865-2874.
- Liu, B., Bai, Q.X., Chen, X.Q., Gao, G.X. & Gu, H.T. (2007) Effect of curcumin on expression of survivin, Bcl-2 and Bax in human multiple myeloma cell line. *Zhongguo shi yan xue ye xue za zhi*, **15**, 762-766.
- Llamas, P., Outeiriño, J., Espinoza, J., Santos, A.B., Román, A. & Tomás, J.F. (2001) Report of three cases of circulating heparin-like anticoagulants. *American Journal of Hematology*, **67**, 256-258.
- Locatelli, F., Aljama, P., Barany, P., Canaud, B., Carrera, F., Eckardt, K.U., Horl, W.H., Macdougall, I.C., Macleod, A., Wiecek, A. & Cameron, S. (2004) Revised European best practice guidelines for the management of anaemia in patients with chronic renal failure. *Nephrology Dialysis Transplantation*, **19 Suppl 2**, ii1-47.
- López, J.A., Kearon, C. & Lee, A.Y. (2004) Deep venous thrombosis. *Hematology American Society Hematology Educational Programme*, 439-456.
- Lötsch, J. & Geisslinger, G. (2006) Current evidence for a genetic modulation of the response to analgesics. *Pain*, **121**, 1-5.
- Mantyh, P.W., Clohisy, D.R., Koltzenburg, M. & Hunt, S.P. (2002) Molecular mechanisms of cancer pain. *Nature Reviews Cancer*, **2**, 201-209.
- Mao, J. (2008) Opioid-induced hyperalgesia. *Pain Clinical Updates*, **XVI**, 1-4.

- Mao, J., Sung, B., Ji, R.R. & Lim, G. (2002) Chronic morphine induces downregulation of spinal glutamate transporters: implications in morphine tolerance and abnormal pain sensitivity. *Journal of Neuroscience*, **22**, 8312-8323.
- Masala, S., Fiori, R., Massari, F., Cantonetti, M., Postorino, M. & Simonetti, G. (2004) Percutaneous kyphoplasty: indications and technique in the treatment of vertebral fractures from myeloma. *Tumori*, **90**, 22-26.
- Mateos, M.V., Oriol, A., Martínez, J., Cibeira, M.T., de Paz, R., Terol, M.J., García-Laraña, J., Bengoechea, E., Martínez, R., Martín, A., de Arriba, F., Palomera, L., Hernández, J., Bello, J.L., Martín, M.L., González, Y., Blade, J., Lahuerta, J.J. & San Miguel, J.F. (2008) Bortezomib (velcade)-melphalan-prednisone (VMP) versus velcade-thalidomide-prednisone (VTP) in elderly untreated multiple myeloma patients: Which is the best partner for velcade: an alkylating or an immunomodulator agent? *Blood (ASH Annual Meeting Abstracts)*, **112**, Abstract 651.
- Mateos, M.V., Garcia-Sanz, R., Colado, E., Olazabal, J. & San-Miguel, J. (2008) Should prophylactic granulocyte-colony stimulating factor be used in multiple myeloma patients developing neutropenia under lenalidomide-based therapy? *British Journal of Haematology*, **140**, 324-326.
- Mavrokokki, T., Cheng, A., Stein, B. & Goss, A. (2007) Nature and frequency of bisphosphonate-associated osteonecrosis of the jaws in Australia. *Journal of Oral and Maxillofacial Surgery*, **65**, 415-423.
- McDonald, R.J., Trout, A.T., Gray, L.A., Dispenzieri, A., Thielen, K.R. & Kallmes, D.F. (2008) Vertebroplasty in multiple myeloma: outcomes in a large patient series. *American Journal of Neuroradiology*, **29**, 642-648.
- Melcher L. (2005) Recommendations for influenza and pneumococcal vaccinations in people receiving chemotherapy. *Clinical Oncology*, **17**, 12-5.
- Menon, S.P., Rajkumar, S.V., Lacy, M., Falco, P. & Palumbo, A. (2008) Thromboembolic events with lenalidomide-based therapy for multiple myeloma. *Cancer*, **112**, 1522-1528.
- Mileshkin, L., Stark, R., Day, B., Seymour, J.F., Zeldis, J.B. & Prince, H.M. (2006) Development of neuropathy in patients with myeloma treated with thalidomide: patterns of occurrence and the role of electrophysiologic monitoring. *Journal of Clinical Oncology*, **24**, 4507-4514.
- Mill, W.B. & Griffith, R. (1980) The role of radiation therapy in the management of plasma cell tumors. *Cancer*, **45**, 647-652.
- Minton, O., Richardson, A., Sharpe, M., Hotopf, M. & Stone, P. (2008) A systematic review and meta-analysis of the pharmacological treatment of cancer-related fatigue. *Journal of the National Cancer Institute*, **100**, 1155-1166.
- Molassiotis, A., Fernandez-Ortega, P., Pud, D., Ozden, G., Platin, N., Hummerston, S., Scott, J.A., Panteli, V., Gudmundsdottir, G., Selvekerova, S., Patiraki, E. & Kearney, N. (2005a) Complementary and alternative medicine use in colorectal cancer patients in seven European countries. *Complementary Therapies in Medicine*, **13**, 251-257.
- Molassiotis, A., Margulies, A., Fernandez-Ortega, P., Pud, D., Panteli, V., Bruyns, I., Scott, J.A., Gudmundsdottir, G., Browall, M., Madsen, E., Ozden, G., Magri, M., Selvekerova, S., Platin, N., Kearney, N. & Patiraki, E. (2005b) Complementary and alternative medicine use in patients with haematological malignancies in Europe. *Complementary Therapies in Clinical Practise*, **11**, 105-110.
- Molassiotis, A., Wilson, B., Blair, S., Howe, T. & Cavet, J. (2010) Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners. *Psychooncology*, Feb 24, epub ahead of print.
- Morgan, G.I., Davies, F.E., Gregory, W.M., Bell, S.E., Szubert, A.J., Cocks, K., Navarro Coy, N., Drayson, M.T., Owen, R.G., Ross, F.M., Jackson, G.H. & Child, J.A. (2009)

The addition of thalidomide to the induction treatment of newly presenting myeloma patients increases the CR rate which is likely to translate into improved PFS and OS. *Blood (ASH Annual Meeting Abstracts)*, **114**, 352.

Mucci-LoRusso, P., Berman, B.S., Silberstein, P.T., Citron, M.L., Bressler, L., Weinstein, S.M., Kaiko, R.F., Buckley, B.J. & Reder, R.F. (1998) Controlled-release oxycodone compared with controlled-release morphine in the treatment of cancer pain: a randomized, double-blind, parallel-group study. *European Journal of Pain*, **2**, 239-249.

Mumford, A.D., O'Donnell, J., Gillmore, J.D., Manning, R.A., Hawkins, P.N. & Laffan, M. (2000) Bleeding symptoms and coagulation abnormalities in 337 patients with AL-amyloidosis. *British Journal of Haematology*, **110**, 454-460.

Murray, S.M. & Pindoria, S. (2009) Nutrition support for bone marrow transplant patients. *Cochrane Database Systemic Reviews*, **21**, CD002920.

Nakazato, T., Ito, K., Ikeda, Y. & Kizaki, M. (2005) Green tea component, catechin, induces apoptosis of human malignant B cells via production of reactive oxygen species. *Clinical Cancer Research*, **11**, 6040-6049.

Nicola, P., Scaramucci, L., Romani, C., Giovannini, M., Maurillo, L., del Poeta, G., Cartoni, C., Arcuri, E., Amadori, S. & De Fabritiis, P. (2006) Opioids in pain management of blood-related malignancies. *Annals of Hematology*, **85**, 489-501.

Nobile-Orazio, E., Barbieri, S., Baldini, L., Marmioli, P., Carpo, M., Premoselli, S., Manfredini, E. & Scarlato, G. (1992) Peripheral neuropathy in monoclonal gammopathy of undetermined significance: prevalence and immunopathogenetic studies. *Acta Neurologica Scandinavica*, **85**, 383-390.

Ossipov, M.H. & Porreca, F. (2005) Challenges in the development of novel treatment strategies for neuropathic pain. *NeuroRx*, **2**, 650-661.

Osterborg, A., Brandberg, Y., Molostova, V., Iosava, G., Abdulkadyrov, K., Hedenus, M. & Messinger, D. (2002) Randomized, double-blind, placebo-controlled trial of recombinant human erythropoietin, epoetin Beta, in hematologic malignancies. *Journal of Clinical Oncology*, **20**, 2486-2494.

Palumbo, A., Bringhen, S., Caravita, T., Merla, E., Capparella, V., Callea, V., Cangialosi, C., Grasso, M., Rossini, F., Galli, M., Catalano, L., Zamagni, E., Petrucci, M.T., De Stefano, V., Ceccarelli, M., Ambrosini, M.T., Avonto, I., Falco, P., Ciccone, G., Liberati, A.M., Musto, P. & Boccadoro, M. (2006) Oral melphalan and prednisone chemotherapy plus thalidomide compared with melphalan and prednisone alone in elderly patients with multiple myeloma: randomised controlled trial. *Lancet*, **367**, 825-831.

Palumbo, A., Bringhen, S., Rossi, D., Magarotto, V., Di Raimondo, F., Ria, R., Offidani, M., Nozzoli, C., Patriarca, F., Callea, V., Benevolo, G., Marasca, R., Guglielmelli, T., Rizzo, M., Grasso, M., Petrucci, M.T., Omede, P., Gaidano, G. & Boccadoro, M. (2008a) A prospective, randomized, phase III study of bortezomib, melphalan, prednisone and thalidomide (VMPT) versus bortezomib, melphalan and prednisone (VMP) in elderly newly diagnosed myeloma patients. *Blood, ASH Annual Meeting Abstracts*, **112**, 652.

Palumbo, A., Facon, T., Sonneveld, P., Bladè, J., Offidani, M., Gay, F., Moreau, P., Waage, A., Spencer, A., Ludwig, H., Boccadoro, M. & Harousseau, J.L. (2008b) Thalidomide for treatment of multiple myeloma: 10 years later. *Blood*, **111**, 3968-3977.

Palumbo, A., Rajkumar, S.V., Dimopoulos, M.A., Richardson, P.G., San Miguel, J., Barlogie, B., Harousseau, J., Zonder, J.A., Cavo, M., Zangari, M., Attal, M., Belch, A., Knop, S., Joshua, D., Sezer, O., Ludwig, H., Vesole, D., Bladé, J., Kyle, R., Westin, J., Weber, D., Bringhen, S., Niesvizky, R., Waage, A., von Lilienfeld-Toal, M., Lonial, S., Morgan, G.J., Orłowski, R.Z., Shimizu, K., Anderson, K.C., Boccadoro, M., Durie, B.G.,

Sonneveld, P. & Hussein, M.A. (2008c) Prevention of thalidomide- and lenalidomide-associated thrombosis in myeloma. *Leukemia*, **22**, 414-423.

Perrone G, Hideshima T, Ikeda H, Okawa Y, Calabrese E, Gorgun G, Santo L, Cirstea D, Raje N, Chauhan D, Baccarani M, Cavo M, Anderson KC (2009). Ascorbic acid inhibits antitumor activity of bortezomib in vivo. *Leukemia*. **23**:1679-86

Peterson, D.E., Bensadoun, R.J. & Roila, F. (2009) Management of oral and gastrointestinal mucositis: ESMO clinical recommendations. *Annals of Oncology*, **20**, 174-177.

Potting, C.M., Uitterhoeve, R., Op Reimer, W.S. & Van Achterberg, T. (2006) The effectiveness of commonly used mouthwashes for the prevention of chemotherapy-induced oral mucositis: a systematic review. *European Journal of Cancer Care (English Language Edition)*, **15**, 431-439.

Rajkumar, S.V., Jacobus, S., Callander, N.S., Fonseca, R., Vesole, D.H., Williams, M.E., Abonour, R., Siegel, D.S., Katz, M. & Greipp, P.R. (2009) Lenalidomide plus high-dose dexamethasone versus lenalidomide plus low-dose dexamethasone as initial therapy for newly diagnosed multiple myeloma: an open-label randomised controlled trial. *Lancet Oncology*, **11**, 29-37.

Ransom, S., Jacobsen, P.B. & Booth-Jones, M. (2006) Validation of the Distress Thermometer with bone marrow transplant patients. *Psychooncology*, **15**, 604-612.

Raphael, J., Ahmedzai, S., Hester, J., Urch, C., Barrie, J., Williams, J., Farquhar-Smith, P., Fallon, M., Hoskin, P., Robb, K., Bennett, M.I., Haines, R., Johnson, M., Bhaskar, A., Chong, S., Duarte, R. & Sparkes, E. (2010a) Cancer pain: part 1: Pathophysiology; oncological, pharmacological, and psychological treatments: a perspective from the British Pain Society endorsed by the UK Association of Palliative Medicine and the Royal College of General Practitioners. *Pain Medicine*, **11**, 742-764.

Raphael, J., Hester, J., Ahmedzai, S., Barrie, J., Farquhar-Smith, P., Williams, J., Urch, C., Bennett, M.I., Robb, K., Simpson, B., Pittler, M., Wider, B., Ewer-Smith, C., Decourcy, J., Young, A., Lioffi, C., McCullough, R., Rajapakse, D., Johnson, M., Duarte, R. & Sparkes, E. (2010b) Cancer Pain: Part 2: Physical, Interventional and Complimentary Therapies; Management in the Community; Acute, Treatment-Related and Complex Cancer Pain: A Perspective from the British Pain Society Endorsed by the UK Association of Palliative Medicine and the Royal College of General Practitioners. *Pain Medicine*, April 26, Epub ahead of print.

Reid, C.M., Martin, R.M., Sterne, J.A., Davies, A.N. & Hanks, G.W. (2006) Oxycodone for cancer-related pain: meta-analysis of randomized controlled trials. *Archives of Internal Medicine*, **166**, 837-843.

Richardson, A. (2006) Sharing the research agenda: confronting what we believe to be the nursing contribution to research in cancer care. *European Journal of Oncology Nursing*, **10**, 91-92.

Richardson, P.G., Barlogie, B., Berenson, J., Singhal, S., Jagannath, S., Irwin, D., Rajkumar, S.V., Srkalovic, G., Alsina, M., Alexanian, R., Siegel, D., Orlowski, R.Z., Kuter, D., Limentani, S.A., Lee, S., Hideshima, T., Esseltine, D.L., Kauffman, M., Adams, J., Schenkein, D.P. & Anderson, K.C. (2003) A phase 2 study of bortezomib in relapsed, refractory myeloma. *New England Journal of Medicine*, **348**, 2609-2617.

Richardson, P.G., Blood, E., Mitsiades, C.S., Jagannath, S., Zeldenrust, S.R., Alsina, M., Schlossman, R.L., Rajkumar, S.V., Desikan, K.R., Hideshima, T., Munshi, N.C., Kelly-Colson, K., Doss, D., McKenney, M.L., Gorelik, S., Warren, D., Freeman, A., Rich, R., Wu, A., Olesnyckyj, M., Wride, K., Dalton, W.S., Zeldis, J., Knight, R., Weller, E. & Anderson, K.C. (2006a) A randomized phase 2 study of lenalidomide therapy for patients with relapsed or relapsed and refractory multiple myeloma. *Blood*, **108**, 3458-3464.

Richardson, P.G., Briemberg, H., Jagannath, S., Wen, P.Y., Barlogie, B., Berenson, J., Singhal, S., Siegel, D.S., Irwin, D., Schuster, M., Srkalovic, G., Alexanian, R., Rajkumar, S.V., Limentani, S., Alsina, M., Orlovski, R.Z., Najarian, K., Esseltine, D., Anderson, K.C. & Amato, A.A. (2006b) Frequency, characteristics, and reversibility of peripheral neuropathy during treatment of advanced multiple myeloma with bortezomib. *Journal of Clinical Oncology*, **24**, 3113-3120.

Rizzo, J.D., Somerfield, M.R., Hagerty, K.L., Seidenfeld, J., Bohlius, J., Bennett, C.L., Cella, D.F., Djulbegovic, B., Goode, M.J., Jakubowski, A.A., Rarick, M.U., Regan, D.H. & Lichtin, A.E. (2008) Use of epoetin and darbepoetin in patients with cancer: 2007 American Society of Clinical Oncology/American Society of Hematology clinical practice guideline update. *Journal of Clinical Oncology*, **26**, 132-149.

Richardson PG, Xie W, Mitsiades C, Chanan-Khan AA, Lonial S, Hassoun H, Avigan DE, Oaklander AL, Kuter DJ, Wen PY, Kesari S, Briemberg HR, Schlossman RL, Munshi NC, Heffner LT, Doss D, Esseltine DL, Weller E, Anderson KC, Amato AA (2009). Single-agent bortezomib in previously untreated multiple myeloma: efficacy, characterization of peripheral neuropathy, and molecular correlations with response and neuropathy. *J Clin Oncol*. **27**(21):3518-25.

Robertson J.D., Nagesh K., Jowitt S.N., Dougal M., Anderson H., Mutton K., Zambon M., Scarffe J.H. (2000) Immunogenicity of vaccination against influenza, Streptococcus pneumoniae and Haemophilus influenzae type B in patients with multiple myeloma. *British Journal of Cancer*, **82**, 1261-1265.

Rowbotham, M.C., Twilling, L., Davies, P.S., Reisner, L., Taylor, K. & Mohr, D. (2003) Oral opioid therapy for chronic peripheral and central neuropathic pain. *New England Journal of Medicine*, **348**, 1223-1232.

Salmon S.E., Samal B.A., Hayes D.M., Hosley H., Miller S.P., Schilling A. (1967) Role of gamma globulin for immunoprophylaxis in multiple myeloma. *New England Journal of Medicine* **277**, 1336-40

Sampson, B.M., Greaves, M., Malia, R.G. & Preston, F.E. (1983) Acquired von Willebrand's disease: demonstration of a circulating inhibitor to the factor VIII complex in four cases. *British Journal of Haematology*, **54**, 233-244.

Sane, D.C., Pizzo, S.V. & Greenberg, C.S. (1989) Elevated urokinase-type plasminogen activator level and bleeding in amyloidosis: case report and literature review. *American Journal of Hematology*, **31**, 53-57.

Sarasquete, M., Gonzalez, M., San Miguel, J. & Garcia-Sanz, R. (2009) Bisphosphonate-related osteonecrosis: genetic and acquired risk factors. *Oral Diseases*, **15**, 382-387.

Savage D.G., Lindenbaum J., Garrett T.J. (1982) Biphasic pattern of bacterial infection in multiple myeloma. *Annals of Internal Medicine*, **96**, 47-50.

Schutter, U., Grunert, S., Meyer, C., Schmidt, T. & Nolte, T. (2010) Innovative pain therapy with a fixed combination of prolonged-release oxycodone/naloxone: a large observational study under conditions of daily practice. *Current Medical Research and Opinion*, **26**, 1377-1387.

Serlin, R.C., Mendoza, T.R., Nakamura, Y., Edwards, K.R. & Cleeland, C.S. (1995) When is cancer pain mild, moderate or severe? Grading pain severity by its interference with function. *Pain*, **61**, 277-284.

Shafqat, A., Einhorn, L.H., Hanna, N., Sledge, G.W., Hanna, A., Juliar, B.E., Monahan, P. & Bhatia, S. (2005) Screening studies for fatigue and laboratory correlates in cancer patients undergoing treatment. *Annals of Oncology*, **16**, 1545-1550.

Shaiova, L., Lapin, J., Manco, L.S., Shasha, D., Hu, K., Harrison, L. & Portenoy, R.K. (2004) Tolerability and effects of two formulations of oral transmucosal fentanyl citrate

- (OTFC; ACTIQ) in patients with radiation-induced oral mucositis. *Supportive Care in Cancer* **12**, 268-273.
- Shammas, M.A., Neri, P., Koley, H., Batchu, R.B., Bertheau, R.C., Munshi, V., Prabhala, R., Fulciniti, M., Tai, Y.T., Treon, S.P., Goyal, R.K., Anderson, K.C. & Munshi, N.C. (2006) Specific killing of multiple myeloma cells by (-)-epigallocatechin-3-gallate extracted from green tea: biologic activity and therapeutic implications. *Blood*, **108**, 2804-2810.
- Shi, H.Y., Mau, L.W., Chang, J.K., Wang, J.W. & Chiu, H.C. (2009a) Responsiveness of the Harris Hip Score and the SF-36: five years after total hip arthroplasty. *Quality of Life Research*, **18**, 1053-1060.
- Shi, Q., Wang, X.S., Mendoza, T.R., Pandya, K.J. & Cleeland, C.S. (2009b) Assessing persistent cancer pain: a comparison of current pain ratings and pain recalled from the past week. *Journal of Pain and Symptom Management*, **37**, 168-174.
- Siemionow, K. & Lieberman, I.H. (2008) Surgical approaches to metastatic spine disease. *Current Opinion in Supportive and Palliative Care*, **2**, 192-196.
- Silberman, J. & Lonial, S. (2008) Review of peripheral neuropathy in plasma cell disorders. *Hematological Oncology*, **26**, 55-65.
- Silverstein, A. & Doniger, D.E. (1963) Neurologic complications of myelomatosis. *Archives of Neurology*, **9**, 534-544.
- Smith, T.J., Coyne, P.J., Staats, P.S., Deer, T., Stearns, L.J., Rauck, R.L., Boortz-Marx, R.L., Buchser, E., Català, E., Bryce, D.A., Cousins, M. & Pool, G.E. (2005) An implantable drug delivery system (IDDS) for refractory cancer pain provides sustained pain control, less drug-related toxicity, and possibly better survival compared with comprehensive medical management (CMM). *Annals of Oncology*, **16**, 825-833.
- Srkalovic, G., Cameron, M.G., Rybicki, L., Deitcher, S.R., Kattke-Marchant, K. & Hussein, M.A. (2004) Monoclonal gammopathy of undetermined significance and multiple myeloma are associated with an increased incidence of venothromboembolic disease. *Cancer*, **101**, 558-666.
- Sung, L., Nathan, P.C., Alibhai, S.M., Tomlinson, G.A. & Beyene, J. (2007) Meta-analysis: effect of prophylactic hematopoietic colony-stimulating factors on mortality and outcomes of infection. *Annals of Internal Medicine*, **147**, 400-411.
- Tassinari, D., Sartori, S., Tamburini, E., Scarpi, E., Tombesi, P., Santelmo, C. & Maltoni, M. (2009) Transdermal fentanyl as a front-line approach to moderate-severe pain: a meta-analysis of randomized clinical trials. *Journal of Palliative Care*, **25**, 172-180.
- Tavares, M. (2003) National Guidelines for the use of complementary therapies in supportive palliative care.
http://www.fih.org.uk/information_library/publications/health_guidelines/complementary.html.
- Thomas, J., Karver, S., Cooney, G.A., Chamberlain, B.H., Watt, C.K., Slatkin, N.E., Stambler, N., Kremer, A.B. & Israel, R.J. (2008) Methylnaltrexone for opioid-induced constipation in advanced illness. *New England Journal of Medicine*, **358**, 2332-2343.
- Thompson, C.A., Kyle, R., Gertz, M., Heit, J., Pruthi, R. & Pardanani, A. (2010) Systemic AL amyloidosis with acquired factor X deficiency: A study of perioperative bleeding risk and treatment outcomes in 60 patients. *American Journal of Hematology*, **85**, 171-173.
- Tovey, P., Broom, A., Chatwin, J., Hafeez, M. & Ahmad, S. (2005) Patient assessment of effectiveness and satisfaction with traditional medicine, globalized complementary and alternative medicines, and allopathic medicines for cancer in Pakistan. *Integrative Cancer Therapies*, **4**, 242-248.

- Trotti, A., Colevas, A.D., Setser, A., Rusch, V., Jaques, D., Budach, V., Langer, C., Murphy, B., Cumberlin, R., Coleman, C.N. & Rubin, P. (2003) CTCAE v3.0: development of a comprehensive grading system for the adverse effects of cancer treatment. *Seminars in Radiation Oncology*, **13**, 176-181.
- Tsavaris, N., Kopterides, P., Kosmas, C., Efthymiou, A., Skopelitis, H., Dimitrakopoulos, A., Pagouni, E., Pikazis, D., Zis, P.V. & Koufos, C. (2008) Gabapentin monotherapy for the treatment of chemotherapy-induced neuropathic pain: a pilot study. *Pain Medicine*, **9**, 1209-1216.
- Twycross, R. & Wilcock, A. (eds.) (2008) *Palliative Care Formulary 3rd edition*. Radcliffe Medical Press, Abingdon.
- Vanstraelen, G., Baron, F., Frere, P., Hafraoui, K., Fillet, G. & Beguin, Y. (2005) Efficacy of recombinant human erythropoietin therapy started one month after autologous peripheral blood stem cell transplantation. *Haematologica*, **90**, 1269-1270.
- Ventafridda, V., Saita, L., Ripamonti, C. & De Conno, F. (1985) WHO guidelines for the use of analgesics in cancer pain. *International Journal of Tissue Reactions*, **7**, 93-96.
- Vriens J, Nilius B, Vennekens R (2008) [Herbal compounds and toxins modulating TRP channels](#). *Curr Neuropharmacol*. **6**:79-96
- Walsh, J.C. (1971) The neuropathy of multiple myeloma. An electrophysiological and histological study. *Archives of Neurology*, **25**, 404-414.
- Wang, M., Weber, D.M., Delasalle, K. & Alexanian, R. (2005) Thalidomide-dexamethasone as primary therapy for advanced multiple myeloma. *American Journal of Hematology*, **79**, 194-197.
- Weber, D.M., Chen, C., Niesvizky, R., Wang, M., Belch, A., Stadtmayer, E.A., Siegel, D., Borrello, I., Rajkumar, S.V., Chanan-Khan, A.A., Lonial, S., Yu, Z., Patin, J., Olesnyckyj, M., Zeldis, J.B. & Knight, R.D. (2007) Lenalidomide plus dexamethasone for relapsed multiple myeloma in North America. *New England Journal of Medicine*, **357**, 2133-2142.
- Weinstein, S.M., Messina, J. & Xie, F. (2009) Fentanyl buccal tablet for the treatment of breakthrough pain in opioid-tolerant patients with chronic cancer pain: A long-term, open-label safety study. *Cancer*, **115**, 2571-2579.
- Weitzman, R., Sauter, N., Eriksen, E.F., Tarassoff, P.G., Lacerna, L.V., Dias, R., Altmeyer, A., Csermak-Renner, K., McGrath, L., Lantwicki, L. & Hohneker, J.A. (2007) Critical review: updated recommendations for the prevention, diagnosis, and treatment of osteonecrosis of the jaw in cancer patients--May 2006. *Critical Reviews in Oncology/Hematology*, **62**, 148-152.
- Wolf, S., Barton, D., Kottschade, L., Grothey, A. & Loprinzi, C. (2008) Chemotherapy-induced peripheral neuropathy: prevention and treatment strategies. *European Journal of Cancer*, **44**, 1507-1515.
- Wu, J.S., Wong, R., Johnston, M., Bezjak, A. & Whelan, T. (1993) Meta-analysis of dose-fractionation radiotherapy trials for the palliation of painful bone metastases. *International Journal of Radiation Oncology, Biology, Physics*, **55**, 594-605.
- Zangari, M., Barlogie, B., Anaissie, E., Saghafifar, F., Eddlemon, P., Jacobson, J., Lee, C.K., Thertulien, R., Talamo, G., Thomas, T., Van Rhee, F., Fassas, A., Fink, L. & Tricot, G. (2004) Deep vein thrombosis in patients with multiple myeloma treated with thalidomide and chemotherapy: effects of prophylactic and therapeutic anticoagulation. *British Journal of Haematology*, **126**, 715-721.
- Zonder, J.A., Barlogie, B., Durie, B.G., McCoy, J., Crowley, J. & Hussein, M.A. (2006) Thrombotic complications in patients with newly diagnosed multiple myeloma treated with lenalidomide and dexamethasone: benefit of aspirin prophylaxis. *Blood*, **108**, 403-404.

